

Mississippi State University

Scholars Junction

College of Agriculture & Life Sciences
Publications and Scholarship

College of Agriculture and Life Sciences

2-24-2014

Quantitative field testing *Heterodera glycines* from metagenomic DNA samples isolated directly from soil under agronomic production.

Yan Li

Gary W. Lawrence

Shien Lu

Mississippi State University, sl332@msstate.edu

Clarissa Balbalian

Mississippi State University, cb237@msstate.edu

Vincent P. Klink

Follow this and additional works at: <https://scholarsjunction.msstate.edu/cals-publications>



Part of the [Genetics Commons](#), and the [Soil Science Commons](#)

Recommended Citation

Li Y, Lawrence GW, Lu S, Balbalian C, Klink VP (2014) Quantitative Field Testing *Heterodera glycines* from Metagenomic DNA Samples Isolated Directly from Soil under Agronomic Production. PLoS ONE 9(2): e89887. <https://doi.org/10.1371/journal.pone.0089887>

This Article is brought to you for free and open access by the College of Agriculture and Life Sciences at Scholars Junction. It has been accepted for inclusion in College of Agriculture & Life Sciences Publications and Scholarship by an authorized administrator of Scholars Junction. For more information, please contact scholcomm@msstate.libanswers.com.

Quantitative Field Testing *Heterodera glycines* from Metagenomic DNA Samples Isolated Directly from Soil under Agronomic Production

Yan Li¹, Gary W. Lawrence¹, Shien Lu¹, Clarissa Balbalian¹, Vincent P. Klink^{2*}

¹ Department of Biochemistry, Molecular Biology, Entomology and Plant Pathology, Mississippi State University, Mississippi State, Mississippi, United States of America,

² Department of Biological Sciences, Mississippi State University, Mississippi State, Mississippi, United States of America

Abstract

A quantitative PCR procedure targeting the *Heterodera glycines* ortholog of the *Caenorhabditis elegans uncoordinated-78* gene was developed. The procedure estimated the quantity of *H. glycines* from metagenomic DNA samples isolated directly from field soil under agronomic production. The estimation of *H. glycines* quantity was determined in soil samples having other soil dwelling plant parasitic nematodes including *Hoplolaimus*, predatory nematodes including *Mononchus*, free-living nematodes and biomass. The methodology provides a framework for molecular diagnostics of nematodes from metagenomic DNA isolated directly from field soil.

Citation: Li Y, Lawrence GW, Lu S, Balbalian C, Klink VP (2014) Quantitative Field Testing *Heterodera glycines* from Metagenomic DNA Samples Isolated Directly from Soil under Agronomic Production. PLoS ONE 9(2): e89887. doi:10.1371/journal.pone.0089887

Editor: Vishal Shah, Dowling College, United States of America

Received: October 5, 2013; **Accepted:** January 27, 2014; **Published:** February 24, 2014

Copyright: © 2014 Li et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The work was supported by the Mississippi Soybean Promotion Board. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: vklink@biology.msstate.edu

Introduction

Plant parasitic nematodes present a major problem for the cultivation of *Glycine max* (soybean) [1–5]. Among them, the dominant problem is the soybean cyst nematode, *Heterodera glycines* [2–5,6]. *H. glycines* was first detected in North Carolina in 1954 [7] and subsequently spread rapidly to as far away as Mississippi by 1957 [8]. The host range of *H. glycines* is broad, updated to over 399 plant species [9–13]. *H. glycines* are a bisexual, cyst-forming species consisting of six life stages including the egg, four juvenile stages (J1–J4), and the adult stage [6,14,15]. The duration of the *H. glycines* life cycle is 3–4 weeks, but this time frame may be influenced by environmental conditions [15]. *H. glycines* feed from their plant host by producing a nurse cell called a syncytium which forms by the fusion and incorporation of approximately 200 cells [16]. Depending upon the environment, several generations of *H. glycines* can be completed in a typical soybean growing season. Adding to management problems, eggs within cysts can remain dormant for 9 years [17]. Management is further complicated because cultivars under production, while experiencing as much as a 30% decrease in yield loss, may appear symptom-free [18]. Infection of soybean by *H. glycines* results in annual losses of \$ 1 billion in yield in the U.S. [19] and about \$15 billion worldwide. To mitigate these losses, management of *H. glycines* is accomplished partly by the use of soybean genotypes that are resistant to infection [20–24]. Although the use of resistant genotypes is a component of recommended control measures [25], consistent use of soybean cultivars with the same sources of resistance can lead to adaptation of the existing *H. glycines* population to cultivars with that source of resistance [26]. Along with the use of resistant

germplasm, the application of crop rotation and nematicides are also used practices in the management of *H. glycines* [27].

A long-used agricultural management strategy is to determine the types and quantities of nematodes present in the soil before planting and this has been particularly important for managing soybean cultivation in *H. glycines*-infested fields [13,28–31]. While these types of tests are traditionally done by trained nematologists, the analyses can be time consuming (i.e. up to 100 days or longer). Therefore, molecular diagnostic tests which could be done in a shorter time frame (i.e. hours) would be useful in the management of many plant-parasitic nematodes [32–39]. Quantitative PCR (qPCR) is a procedure that specifically assays the PCR reaction by incorporating a molecular tag whose amplification products are measured during the PCR reaction. The qPCR method works quantitatively in pathogen detection and measurement because it detects the pathogen by the using the amount of DNA present in a sample to obtain a cycle threshold (Ct) value which corresponds to the level of infection [40]. The qPCR methodology has been shown to be a very powerful tool in pathogen detection, used to identify and quantify various bacteria, fungi, and viruses [41–44]. Population estimates for nematode species such as *M. javanica*, *Pratylenchus zaeae*, and *Xiphinema elongatum* have been determined with qPCR on native soil samples from trial plots [45]. The qPCR procedure has also been used to determine the quantity of the plant parasitic nematode *Rotylenchulus reniformis* from metagenomic DNA isolated directly from field plots [46].

It is hypothesized that conserved gene sequences can be used in determining the number of *H. glycines* from metagenomic DNA isolated directly from soil. In the analysis presented here, highly conserved *H. glycines* homologs of the *C. elegans uncoordinated* gene family genes [47] were screened for their use as biomarkers against

Table 1. PCR primer pairs for *H. glycines* homologs of *uncoordinated* (*unc*) class genes.

<i>H.glycines</i> gene	<i>H.glycines</i> est	Product Size	Primer	TM
Hg-unc-9	CB281382	124	F-CCATGGTGCGCTATTTGTCA	62.95
			R-CTGCCCCAAATTGTTTGAA	63
Hg-unc-9	CB281382	253	F-GCGCGAAAGGACGATGATTTTTTC	66.5
			R-CTTACGGCCGGACGAATACCTCTC	66.6
Hg-unc-22	CB378705	120	F-GACGAAATTGTGGCCGAGTC	62.91
			R-AAATTGTCCCGCTCTCTT	63.11
Hg-unc-31	CB378080	110	F-CTCCGATGGTTGTCGCTAC	62.91
			R-GGTTGAGCAACCGTCTTTGC	63.05
Hg-unc-52	CK350534	114	F-ACCGCAGGTGTACGATGGTT	62.69
			R-CCGTAGGCGGTCACTTTGTC	62.92
Hg-unc-78	CB238521	97	F-CGTTTTTGGGACACCACACA	62.79
			R-TGCTGTCTCTCAGACCAGAA	63.04
			qPCR probe GAAGTCGGAGTTCGCTCTCTTTTCG	70.98
Hg-unc-78	CB238521	313	F-GTGAGACCAATCGGGCAAATC	66.4
			R-GAAAGGAGGGCCTTCGAAATGG	66.5
Hg-unc-89	CB379143	120	F-GCGCGGTACTGACGAAAGTC	63.19
			R-GCAGGACAGTTCCGCATTC	63.04
Hg-unc-89	CB379143	289	F-CCCCTACACACATTTCCGCAGTC	66.3
			R-CAGCCGACCATCGAGTCCATAC	66.5
Hg-unc-101	CB379764	94	F-CATGCAAGGCAACAGATTCG	62.7
			R-TAACAGCGCACATCCAACG	63.09
Hg-unc-115	CK350435	103	F-ACGGAAGTCGCGTATTCAA	63.06
			R-GTCGTTGTCCACGGAAGAGG	63.01
Hg-dys-1	CB934909	124	F-GCTATTTGCCGTCGAACAA	63.28
			R-TTGTCCTCAATCTCGGGCTAT	62.92
Hg-dys-1	CB934909	303	F-GTTCCGATCGTTGGACTTCG	66.5
			R-GCTGGTGCATTGCCTCTGTTTC	66.4
Hg-nep-1	CB824545	114	F-TATTCGGGCGTCAAAAATGC	63.04
			R-GCCAATCACTGCTCCAATCC	62.9
Hg-MRCK-1 (control)	CB380016	125	F-CCACCACACGTCCAATTT	63.27
			R-GAAGGTGAAGCCGATGAACG	63.02

doi:10.1371/journal.pone.0089887.t001

other phylogenetically related and unrelated nematodes [48]. The molecular diagnostic was able to specifically identify *H. glycines* from metagenomic DNA samples isolated directly from soil under agricultural production. Furthermore, a single, *H. glycines* could be identified from the metagenomic DNA sample. The methodology is rapid, and can be performed *en masse* in a single afternoon.

Results

Probe Identification

Alkharouf et al. [49] analyzed 8,334 *H. glycines* unigenes that resulted in the identification of numerous genes involved in essential aspects of their biology. From those sequences, 11 genes were targeted for their use as molecular probes to determine the level of soil infestation by *H. glycines* (Table 1). The 11 primer pairs, designed from these gene sequences, were used in a series of PCR reactions to identify genes that would robustly amplify *H.*

glycines DNA isolated from pure single cysts as compared to a control gene (Figure 1). As demonstrated by an increase in the amount of amplification product, semiquantitative PCR tests demonstrated Hg-unc-78 primer pairs were able to detect increasing amounts of DNA from increasing numbers of *H. glycines* J2 (Figure 2). Gene sequencing confirmed the PCR amplicon was Hg-unc-78 (data not presented).

Detecting *H. glycines* using Metagenomic DNA Isolated Directly from Soil

The experiments then focused on determining if *H. glycines* DNA could be isolated and detected directly from soil samples. Metagenomic DNA isolation, followed by PCR, demonstrated that the Hg-unc 78 amplification product could be detected as a band in samples with as few as 1 added J2 (Figure 3). To further confirm the specificity of Hg-unc 78 primers, experiments on agriculturally significant off-target plant-parasitic nematodes




Figure 1. Screening of Hg-unc primer pairs. Lane 1, Hg-unc 9; Lane 2, Hg-unc 22; Lane 3, Hg-unc 31; Lane 4, Hg-unc 52; Lane 5, Hg-unc 115; Lane 6, Hg-unc 101; Lane 7, Hg-dys-1; Lane 8, Hg-nep-1; Lane 9, Hg-unc 89; Lane 10, Hg-unc 78; Lane 11, Hg-MRCK-1. doi:10.1371/journal.pone.0089887.g001

including *R. reniformis* [50] (Hoplolaimidae) and *M. incognita* (root knot nematodes, RKN) (Heteroderidae), along with *H. glycines*, DNA were used in PCR experiments. No amplification was achieved in reactions containing *R. reniformis* or *M. incognita* DNA, but amplification was successful on reactions containing *H. glycines* DNA (Figure 4). This outcome indicates that the primers designed to amplify Hg-unc 78 is specific to *H. glycines*. Since it was possible to amplify target *H. glycines* DNA from soil containing as little as a single *H. glycines* J2 nematode, experiments were done whereby metagenomic DNA was isolated directly from field samples containing numerous parasitic and free-living nematodes as well as other living organisms. Soil samples were collected in triplicate and a visual assessment of the nematodes present within the soil sample were determined (Table 2). This procedure determined that the soil contained other species of nematodes that are more distantly related to *H. glycines*, including *Hoplolaimus* (sp.) [51] (Hoplolaimidae) and *Mononchus* (sp.) [52] (Mononchidae). Known numbers of *H. glycines* were then added to the soil samples and the metagenomic DNA was isolated directly from the soil as described previously. The amplification profile of the Hg-unc-78 DNA is shown (Figure 5). The demonstration of increasing amounts of amplification product in samples having increasing amount of added *H. glycines* indicated that it would be possible to adapt qPCR to the detection method.




Figure 2. DNA amplification of Hg-unc-78 from different numbers of pure, extracted *H. glycines* (SCN) J2 cultured in soil in greenhouse. Lanes 1–3, 1 SCN; Lanes 4–6, 10 SCN; Lanes 7–9, 100 SCN; Lanes 10–12, 1,000 SCN. doi:10.1371/journal.pone.0089887.g002




Figure 3. DNA amplification of Hg-unc-78 from metagenomic DNA isolated directly from soil containing different numbers of *H. glycines* (SCN). Lanes 1–3, 1 SCN; Lanes 4–6, 10 SCN; Lanes 7–9, 100 SCN; Lanes 10–12, 1,000 SCN; Lane 13, without primer (control); Lane 14, without DNA (control). doi:10.1371/journal.pone.0089887.g003

qPCR Estimate of *H. glycines*

The qPCR experiments began with pure, greenhouse cultured *H. glycines* J2s (Figure 6). From this standard curve, it is demonstrated that the Hg-unc-78 qPCR primers are able to measure, quantitatively, the different amount of *H. glycines* DNA within the samples. This measurement allows for the correlation of the quantity of the amplified DNA amount to estimate number of *H. glycines* J2s. The same procedure then was used to estimate the number of *H. glycines* J2s directly from metagenomic DNA isolated directly from soil samples (Figure 7). Subsequently, actual soil samples under agricultural production were assayed using qPCR for *H. glycines* (Table 3). Metagenomic DNA was isolated from those samples and used in qPCR experiments. Since the number of *H. glycines* is a pool of both cysts and J2s, it was necessary to estimate the number of J2s per cyst. Using that estimate, a correlation between the counted *H. glycines* as compared to the qPCR estimate is provided (Table 3). DNA sequencing of the qPCR product demonstrated the specificity of the reaction by revealing the amplification product was Hg-unc-78 (data not presented).

Discussion

The determination of the numbers of specific types of plant parasitic nematodes in the soil is important in agriculture because it allows for understanding threshold populations that could detrimentally affect plant productivity. This knowledge, in turn, will allow growers to make informed choices in terms of cultivar and/or crop selection as nematode populations can vary from year to year. The analysis presented here resulted in the development of a simple and reliable molecular diagnostic technique that determines the number of *H. glycines* from metagenomic DNA isolated directly from field soil under agricultural production. During the course of the analysis, gene sequences for *H. glycines* homologs of *C. elegans unc* genes were identified, tested and analyzed for their reliability in molecular diagnostics. During the analysis, qPCR primers were designed and used on metagenomic



Figure 4. The specificity of the Hg-unc78 PCR reaction from DNA isolated from different numbers of pure, extracted off-target nematodes as compared to DNA isolated from pure, extracted *H. glycines*. Lanes 1–3; 100 Rr; Lanes 4–6, 1,000 Rr; Lanes 7–9, 100 RKN; Lanes 10–12, 1,000 RKN; Lanes 13–15, 100 SCN; Lanes 16–18, 1,000 SCN; Lane 19, without primers (control); Lane 20, without DNA (control). Rr, *Rotylenchulus reniformis*; RKN, *Meloidogyne incognita*. doi:10.1371/journal.pone.0089887.g004

Table 2. Quantification of major nematode fauna from field soil.

Microplots	<i>Hoplolaimus</i> spp	<i>Mononchus</i> spp	Free living
1	1287.5	772.5	772.5
2	772.5	2317.5	257.5
3	257.5	0	515
4	257.5	257.5	515

doi:10.1371/journal.pone.0089887.t002

DNA isolated directly from actual soil samples from real agricultural sites. While the focus of the analysis was low numbers of *H. glycinis* (1–1,000 individual J2s), it is expected that the methodology would also be reliable for analyzing field samples that are more highly infested.

Hg-unc-78 as a Molecular Probe

Gene sequences analyzed in Alkharouf et al. [49] was mined for genes that could be used as reliable molecular probes for the presence of *H. glycinis* in actual field samples under agricultural production. Genes that function in nematode movement, an important aspect of parasitism, and were also conserved in their primary structure to *C. elegans* were identified and used for PCR primer design. Many of these genes were *unc* genes, first described in *C. elegans* [53]. The experiments presented here show that it is possible to use highly conserved DNA sequences that are homologous to *unc* genes from *H. glycinis* and use them to specifically amplify gene fragments from as few as 1 J2. The specificity was shown when attempts in amplification reactions of off-target nematodes and, in particular, metagenomic DNA isolated directly from soil samples. DNA sequence analyses of the DNA amplification products from the PCR reactions show that the amplification product perfectly matches its target DNA from *H. glycinis*. Lastly, using the same primers to amplify target DNA from metagenomic DNA isolated directly from soil samples reveals that it is possible to amplify the target as shown by sequencing reactions of that amplification product to as low as a single *H. glycinis* J2. Furthermore, in regard to the major nematode species found in Mississippi soils, the procedure appears to distinguish *H. glycinis*. However, all nematodes have not been examined.

Quantitative PCR

The qPCR experiments were designed with the goal of determining the numbers of *H. glycinis* in soil samples collected from actual field sites under agricultural production. From the described PCR studies, it was determined that it would be possible




Figure 5. DNA amplification of the Hg-unc78 from metagenomic DNA samples. Lanes 1–3, metagenomic DNA including 1 SCN J2; Lanes 4–6, metagenomic DNA including 10 SCN J2; Lanes 7–9, metagenomic DNA including 100 SCN J2; Lanes 10–12, metagenomic DNA including 1,000 SCN J2; Lane 13, cloned gene without primers (control); Lane 14, without DNA (control); Lane 15, positive control, amplification product is Hg-unc 78 amplified from the cloned gene. doi:10.1371/journal.pone.0089887.g005

to identify a single *H. glycinis* from metagenomic DNA. Similar observations have been made for *R. reniformis* [46]. The DNA samples isolated from pure cultures of *H. glycinis* could then be used in the development of a standard curve from which it was possible to estimate the number of *H. glycinis*. With this capability, it was then tested whether it would be possible to take metagenomic DNA samples isolated from soil having predefined numbers of *H. glycinis* to quantify the number of *H. glycinis*. As shown, the Hg-unc-78 primer set could be used in experiments to accurately estimate the number of J2s in the sample. The quantitative analysis, using qPCR, demonstrates that low numbers of *H. glycinis* can be identified reliably from the metagenomic samples. Notably, the quantity of *H. glycinis* has been determined through visual analysis from actual soil samples under agricultural production, serving as experimental samples. Metagenomic DNA then was isolated from those samples and used for subsequent qPCR analyses. The analysis demonstrated that the qPCR did not perfectly determine the exact number of *H. glycinis* in the agricultural samples. The same observation was made for *R. reniformis* [46]. However, it is well known that *H. glycinis* are not evenly distributed within soil samples [27]. Thus, the limits of detection may have been met. Further experimentation is required to explore this problem further.

A key question remaining is whether different compositions of soil will affect the DNA isolation efficiency. In Mississippi where the experiments were carried out, there are 5 major soil types. The soil types vary among alfisols, entisols, inceptisols, ultisols and vertisols [54]. It is possible that these different soil compositions could affect the efficiency of the DNA isolation and amplification. While no major influence was observed on amplification from the different sample types used in the various analyses, a thorough analysis on different soil types from the obtained unknown samples requires further testing.

Materials and Methods

Ethics Statement

Since the focal point of the research was plant parasitic nematodes, IACUC regulations were not relevant to this study.

Materials Statement

No specific permits were required for the described field studies because the soil samples were collected on the grounds owned by Mississippi State University and contained no endangered or protected species as determined by a stereomicroscopic analysis of soil.

Plant Inoculation

Prior to nematode inoculation, *G. max* seedlings are grown in a sterilized sand: soil mix (1:1) in clay pots for a period of 7 to 14 days. The nematodes of second stage juveniles were counted, concentrated and diluted to a final concentration of 2,000 J2 per 3 ml which were then added to each root. This meant that 3 ml of inoculum that contained 2,000 nematodes were added to each root system on each plant by inoculating on two spots evenly.

Plant and Nematode Culture

The target nematode *H. glycinis* and off-target *M. incognita* (root knot nematode, RKN) [55–56] (Heteroderidae) and *R. reniformis* (reniform nematode) [48] (Hoplolaimidae) were cultured under ambient conditions in a greenhouse at the Mississippi Agriculture and Forestry Experiment Station (MAFES), RR Foil Plant Science Research Center, North Farm, Mississippi State University. Supplemental fluorescent light was provided to bring the day




Figure 6. Estimation of *H. glycines* population size by qPCR. Standard curve obtained using pure greenhouse samples, (b) soil samples. The statistical significance between the actual number of SCN and the estimated number are provided. doi:10.1371/journal.pone.0089887.g006

length to a 16 hour day/8 hour night cycle. Temperatures were kept in a constant temperature range between 28.9–34.4°C. Nematodes were cultured in 500 cm³ diameter clay pots for a period of 2–6 months in a 50:50 mixture of a fine sandy loam (46.25% sand, 46.50% silt, and 7.25% clay) and sand (93.00% sand, 5.75% silt, and 1.25% clay). Harvesting cysts was accomplished by massaging the infected roots in water. Massaging was achieved by placing the infected root mass between the index finger and thumb. The index finger and thumb were gently rubbed together with the root mass between them. This gentle rubbing activity dislodged the cysts so that they could be collected on a 150 µm pore sieve under a constant water steam. The cysts are not damaged because of their protective hardened nature. Cysts were collected by rinsing them through nested 850 µm pore sieve (debris) and a 150 µm sieve (cyst). Then cysts were rinsed out of the 150 µm pore sieve into a beaker with 100 ml of water. Juveniles were collected by rinsing them through 250 µm pore sieve (debris) and 45 µm pore sieve (J2) and transferred to a beaker with 100 ml water in it for counting. Cysts were crushed by using Janke & Kunkel IKA-WERK crushing machine (IKA-WORKS,

INC), for 30 seconds to 1 minute with a timer. Removal of debris smaller than the eggs was done by washing the slurry through a 25 µm pore sieve. The eggs were transferred to a beaker with 100 ml water in it for counting [57].

Soybean Cyst Nematode DNA Isolation

To isolate DNA from a single cyst, a single cyst was picked up under a stereomicroscope and placed into a 1.5 ml microcentrifuge tube, with 14 µl sterile distilled water. The tube was placed in –20°C for 1 hour to freeze the cyst (or liquid nitrogen for 1 min). A glass rod sterilized with 75% ethanol was used to grind the cyst containing mixture until it is thawed. A 3 µl 10×PCR buffer and 3 µl proteinase K was then added. The material was incubated at –20°C for at least 2 hours. The solution was incubated for 90 min at 65°C to denature the protein, and 15 min at 9°C to unwind the DNA. The solution was centrifuged for 1 min at 8,000 rpm/h with the supernatant transferred to a new 1.5 ml microcentrifuge tube. The DNA was stored at –20°C for later PCR analysis.




Figure 7. Estimation of *H. glycines* population size by qPCR. Standard curve obtained from metagenomic DNA isolated directly from soil samples. The statistical significance between the actual number of SCN and the estimated number are provided. doi:10.1371/journal.pone.0089887.g007

Table 3. qPCR estimation of *H. glycines* from field soil samples.

Sample	J2/pint	Cysts/pint	Cyst to J2*	Actual NO. J2/pint	qPCR J2 estimate/pint	fold difference
1	0	8	1,600	1,600	8,394.38921	0.19
2	0	16	3,200	3,200	5,599.6761	0.57
3	0	24	4,800	4,800	7,692.002288	0.62
4	24	16	3,200	3,224	4,928.816234	0.65
5	0	16	3,200	3,200	2,369.447	1.35
6	63	24	4,800	4,863	1,857.50756	2.62
7	8	8	1,600	1,608	421.9439991	3.81
8	0	8	1,600	1,600	370.9181429	4.32
9	16	47	9,400	9,416	1,557.579433	6.05
10	32	32	6,400	6,432	747.7392119	8.62

doi:10.1371/journal.pone.0089887.t003

Gene Screening

H. glycines genes identified in Alkharouf et al. [49] were used for primer design according to Showmaker et al. [46] who used the identical methodology for quantitative molecular detection of *Rotylenchulus reniformis* DNA from metagenomic samples isolated from soil. Highly conserved genes from Group 1 [49], having 266 genes with *E*-values between 0 and $1E-100$, were the focus of the analysis because no other *Heterodera* spp. agricultural important species has been published to be found in Mississippi soils at this time. Many of these genes included the *uncoordinated* (*unc*) class were first described in *C. elegans*. The genetically-defined *unc* genes group contains 114 different members and function in muscle focal adhesion, architecture and stimulation [53].

Metagenomic DNA Isolation

Metagenomic DNA from soil isolations were conducted by using the Powersoil DNA extraction kit (MO BIO Laboratories, Inc; Carlsbad, CA) using the manufacturer's protocol with modifications. The modifications included using 0.3 grams of soil instead of 0.25 grams of soil in step for developing the quantitative PCR standard curve and adding 0.1 ml of nematode suspension (water containing the numbers of J2 extracted from greenhouse cultured soil samples). In its place, 0.1 ml of the nematode suspension, extracted from either greenhouse pots or field soil, was pipetted into the bead beating tube. Secondly, in the step when first instructed to remove the supernatant, a standard volume of 550 ml of supernatant was removed from each tube. This ensured for downstream applications. The DNA was eluted from the spin column in 50 μ l of nuclease free water (Promega; Madison, WI) and stored at -20°C .

PCR Reaction Conditions

For PCR, a 50 μ l PCR reaction consisting of 2 μ l DNA template, 5.0 μ l $10\times$ PCR buffer, 1.0 μ l dNTP, 3.0 μ l MgCl_2 , 0.25 μ l recombinant Taq Polymerase, 1.25 μ l of 100 nM forward and reverse primers each, 36. μ l nuclease free water (Ambion) (Promega) was used. The reaction conditions, as reported by Agudelo et al. [39] were modified to include a 2 minute pre-denaturation step at 94°C . The procedure then followed the Agudelo et al. [39] protocol that included a denaturation at 94°C for 45 sec, annealing at 59°C for 45 sec and primer extension at 72°C for 60 sec for 40 cycles. The PCR reaction products were run out by gel electrophoresis on a 1% agarose gel with 0.01% SYBR-Green incorporated into the gel. The DNA amplification

products were visualized and recorded with digital imagery using a FOTO/Analyst Apprentice System (FOTODYNE Inc.; Hartland, WI).

Cloning and Sequencing

After using Accuprime polymerase for the PCR, the DNA amplification products were extracted on the gel by using the gel extraction kit (QIAQuick Gel Extraction Kit), mixing 4 μ l of the gel purified DNA with 1 μ l of salt solution and 1 μ l of TOPO vector for the TOPO cloning reaction (pENTR Directional TOPO Cloning Kits, Invitrogen) in a 1.5 ml microcentrifuge tube. The tube was incubated at room temperature for 5 min and then placed on ice. Chemically competent *E. coli* stored at -80°C was thawed on ice. Transformation began by distributing the *E. coli* cells in a separate 1.5 ml microcentrifuge tube, at a volume of 20 μ l each, which is proceeded on ice. Then, 3 μ l of TOPO cloning reaction was added to the *E. coli* cells, tapping the tube gently. The *E. coli* solution were incubated on ice for 15 min followed by heat shocking the *E. coli* cells at 42°C for 30 seconds on the heat block. The *E. coli* solution then was placed back on ice immediately. Subsequently, 100 μ l of SOC media was added to the tube and shaken at 37°C for 1 hour. After incubation, 70 μ l of the solution were plated out on LB-kanamycin plates. The plates were incubated at 37°C overnight for 12–24 hours. The next day, single colonies were picked and placed in a 50 ml centrifuge tube containing LB-kanamycin culture agar. The tubes were cultured for 37°C for no more than 16 hours, usually around 15 hours. A plasmid prep was done on the culture (QIAPrep, Miniprep, QIAGEN).

To confirm that the DNA amplification in both PCR and qPCR reactions were products of *H. glycines* DNA and not spurious amplification of off-target DNA, DNA amplification products were run out electrophoretically on and then isolated from the 1% agarose gels. The DNA was purified using the QIAquick Gel Extraction Kit (QIAGEN; Valencia, CA) as according to the manufacturer's specifications. The isolated DNA was ligated into the pGEM-T Vector System II (Promega). The ligation reaction was shuttled into competent JM109 *E. coli* cells and selected on 50 $\mu\text{g}/\text{ml}$ ampicillin on LB-agar plates. Colonies were selected and grown in liquid culture in LB media containing 50 $\mu\text{g}/\text{ml}$ ampicillin. Plasmid DNA was isolated from the bacteria using the QIAPrep kit (QIAGEN). The DNA from the plasmid preps was sequenced to determine if the DNA amplification product was correctly amplifying the proper target. The DNA sequence was trimmed using the Crimson Editing freeware (<http://www.>

crimsoneditor.com/). In this procedure, the pGEM-T Vector DNA sequence was trimmed leaving the qPCR-generated sequence. The trimmed sequence was blasted in GenBank using the blastn query option. This additional quality control step demonstrated the accuracy of the qPCR reaction conditions.

Quantitative PCR

Taqman 6-carboxyfluorescein (6-FAM) qPCR probes (MWG Operon; Birmingham, AL) were used. The 6-FAM probes have a maximum excitation at 495 nm and maximum emission at 520 nm. The qPCR quencher was the Black Hole Quencher (BHQ1) (MWG Operon), with maximum excitation at 534 nm. Assays were conducted for primers that produced a single amplicon and had no off target amplification. The qPCR reaction conditions included a 20 μ l Taqman Gene Expression Master Mix (Applied Biosystems; Foster City, CA), 0.9 μ l 100 μ M forward primer, 0.9 μ l 100 μ M reverse primer, 2 μ l 2.5 μ M 6-FAM (MWG Operon) probe and 4.4 μ l metagenomic template DNA. The conditions were a denaturation at 94°C for 45 sec, annealing at 54°C for 45 sec and primer extension at 72°C for 60 sec for 40 cycles. The qPCR reactions were performed on an ABI 7300 (Applied Biosystems).

To generate a standard curve for the amount of *H. glycines* in a soil sample, estimates of approximately 1,000 nematodes in 0.1 ml of water were placed into the Powersoil DNA isolation kit bead beating tubes and extracted as described previously, replicated this by 2 more times. A 1:10 serial dilution series of DNA extracted from approximately 1,000 nematodes was created and used for generation of the standard curve by qPCR. To evaluate the accuracy of the standard curve, samples containing 0, 1, 10 and 100 *H. glycines* nematodes were generated by carefully hand

collecting them under a stereoscope, 3 replicates of each, and isolated the DNA by the Powersoil DNA isolation kit as described previously. The qPCR methodology works quantitatively because it detects pathogens by using the amount of DNA present in a sample to obtain a cycle threshold (C(t)) value which corresponds to the amount of target DNA (Livak and Schmittgen, 2001). The lower the C(t), the greater the amount of the corresponding DNA (target organism) is present in a sample.

H. glycines Estimation

50 g of soil was weighed and put into a cylinder. Subsequently, 50 ml of water was put in the cylinder. Because of the volume of the soil, the total volume increased to 70 ml, meaning the 50 g of soil was equal to a volume of 20 ml. With 50 g taking up 20 ml, 0.3 g of soil takes $(0.3 \text{ g} \times 20 \text{ ml}) / 50 \text{ g} = 0.12 \text{ ml}$. So the estimated number of J2 in 1 pint = $473.176 \text{ ml} \times C(t) / 0.12$.

Acknowledgments

The authors thank Dr. Larry Heatherly and the Mississippi Soybean Promotion Board and the Mississippi Agricultural and Forestry Experimental Station (MAFES). Dr. Scott Willard, Department of Biochemistry, Molecular Biology, Entomology and Plant Pathology (BMPEP), Mississippi State University provided greenhouse space. Yixiu Pinnix (BMPEP) provided technical support. The authors are thankful to Dr. Tom Allen, Delta Research and Extension Center, Stoneville, MS.

Author Contributions

Conceived and designed the experiments: YL GL VK. Performed the experiments: YL. Analyzed the data: YL GL SL CB VK. Contributed reagents/materials/analysis tools: YL GL CB VK. Wrote the paper: YL GL VK.

References

- Wrather JA, Chambers AY, Fox JA, Moore WF, Sciombato GL (1995) Soybean disease loss estimates for the southern United States, 1974–1994. *Plant Disease* 79: 1076–1079.
- Wrather JA, Anderson TR, Arsyad DM, Tan Y, Ploper LD, et al. (2001a) Soybean disease loss estimates for the top ten soybean producing countries in 1998. *Canadian Journal of Plant Pathology* 23: 115–121.
- Wrather JA, Steinstra WC, Koenning SR (2001b) Soybean disease loss estimates for the United States from 1996–1998. *Canadian Journal of Plant Pathology* 23: 122–131.
- Wrather JA, Koenning SR, Anderson TR (2003) Effect of diseases on soybean yields in the United States and Ontario (1999–2002). Online. *Plant Health Progress* doi: 10.1094/PHP-2003-0325-01-RV. <http://www.plantmanagement.network.org/sub/php/review/2003/soybean/> > (19 November 2003).
- Wrather JA, Koenning SR (2006) Estimates of disease effects on soybean yields in the United States 2003–2005. *Journal of Nematology* 38: 173–180.
- Ichinohe M (1952) On the soybean nematode, *Heterodera glycines* n. sp., from Japan. *Magazine of Applied Zoology* 17: 1–4.
- Winstead NN, Skotland CB, Sasser JN (1955) Soybean cyst nematodes in North Carolina. *Plant Disease Reporter* 39: 9–11.
- Spears JF (1957) Review of soybean cyst nematode situation for presentation at public hearing on the need for Federal Domestic Plant Quarantine, July 24, 1957.
- Epps JM, Chambers AY (1958) New host records for *Heterodera glycines* including one in the Labiate. *Plant Disease Reporter* 42: 194.
- Riggs RD, Hamblen ML (1962) Soybean-cyst nematode host studies in the Leguminosae. *Arkansas Agriculture Experimental Station Reporter Series* 110 Fayetteville AR 17p.
- Riggs RD, Hamblen ML (1966a) Additional weed hosts of *Heterodera glycines*. *Plant Disease Reporter* 50: 15–16.
- Riggs RD, Hamblen ML (1966b) Further studies on the host range of the soybean-cyst nematode. *Arkansas Agriculture Experimental Station Bulletin* 718 Fayetteville AR 19p.
- Niblack TL, Arelli PR, Noel GR, Opperman CH, Orf JH, et al. (2002) A revised classification scheme for genetically diverse populations of *Heterodera glycines*. *Journal of Nematology* 34: 279–288.
- Lauritis JA, Rebois RV, Grancy LS (1983) Development of *Heterodera glycines* Ichinohe on soybean, *Glycine max*. *Journal of Nematology* 15: 272–280.
- Davis EL, Tylka GL (2000) Soybean cyst nematode disease. *The Plant Health Instructor*. DOI: 10.1094/PHI-I-2000-0725-01.
- Jones MGK, Northcote DH (1972) Nematode-induced syncytium—a multinucleate transfer cell. *J Cell Sci* 10: 789–809.
- Inagaki H, Tsutsumi M (1971) Survival of the soybean cyst nematode, *Heterodera glycines* Ichinohe (Tylenchida: Heteroderidae) under certain storage conditions. *Applied Entomology and Zoology (Jpn)* 8: 53–63.
- Wang J, Meyers D, Yan Y, Baum T, Smant G, et al. (2003) The soybean cyst nematode reduces soybean yield without causing obvious symptoms. *Plant Disease* 87: 623–628.
- Lawton K (2010) Defeating soybean cyst nematode requires planting non-host crops and varieties with different sources of resistance. *The Progressive Farmer*: 28.
- Ross JP, Brim CA (1957) Resistance of soybeans to the soybean cyst nematode as determined by a double-row method. *Plant Dis Rep* 41: 923–924.
- Ross JP (1958) Host-Parasite relationship of the soybean cyst nematode in resistant soybean roots. *Phytopathology* 48: 578–579.
- Epps JM, Hartwig EE (1972) Reaction of soybean varieties and strains to soybean cyst nematode. *Journal of Nematology* 4: 222.
- Concibido VC, Diers BW, Arelli PR (2004) A decade of QTL mapping for cyst nematode resistance in soybean. *Crop Sci* 44: 1121–1131.
- Shannon JG, Arelli PR, Young LD (2004) Breeding for resistance and tolerance. Pp. 155–180 in D. P. Schmitt, J. A. Wrather, and R. D. Riggs, eds. *Biology and management of soybean cyst nematode*, 2nd ed. Marceline, MO: Schmitt & Associates of Marceline.
- Riggs RD, Niblack TL (1999) Soybean cyst nematode. In: Hartman GL, Sinclair JB, Rupe JC, editors. *Compendium of Soybean Diseases*. Fourth Edition. St. Paul: APS Press; 52–53.
- Luedders VD, Anand SC (1989) Attempt to select a cyst nematode population on soybean plant introduction 437654. *Journal of Nematology* 21: 264–267.
- Niblack TL (2005) Soybean cyst nematode management reconsidered. *Plant Disease* Vol. 89 No.10.
- Cobb NA (1918) Estimating the Nema population of the soil. *Agricultural Technology Circular Bureau of Plant Industry, US Department of Agriculture*, 1: 1–48.
- Golden AM, Epps JM, Riggs RD, Duclos LA, Fox JA, et al. (1970) Terminology and identity of infraspecific forms of the soybean cyst nematode (*Heterodera glycines*). *Plant Dis Rep* 54: 544–546.
- Riggs RD, Schmitt DP (1988) Complete characterization of the race scheme for *Heterodera glycines*. *Journal of Nematology* 20: 392–395.
- Riggs RD, Schmitt DP (1991) Optimization of the *Heterodera glycines* race test procedure. *Journal of Nematology* 23: 149–154.

32. Harris TS, Sandall LJ, Powers TO (1990) Identification of single Meloidogyne juveniles by polymerase chain reaction amplification of mitochondrial DNA. *Journal of Nematology* 22: 518–524.
33. Roosien J, Van Zandvoort PM, Folkertsma RT, Van der Voort JN, Goverse A, et al. (1993) Single juveniles of the potato cyst nematodes *Globodera rostochiensis* and *G. pallida* differentiated by randomly amplified polymorphic DNA. *Parasitology* 107: 567–572.
34. Gasser RB, Monti JR (1997) Identification of parasitic nematodes by PCR-SSCP of ITS-2 rDNA. *Molecular Cell Probes* 11: 201–209.
35. Powers TO, Harris TS (1993) A polymerase chain reaction method for identification of five major Meloidogyne species. *Journal of Nematology* 25: 1–6.
36. Cenis JL (1993) Identification of four major *Meloidogyne* spp. by random amplified polymorphic DNA (RAPD-PCR). *Phytopathology* 83: 76–80.
37. Semblat JP, Wajnberg E, Dalmasso A, Abad P, Castagnone-Sereno P (1998) High-resolution DNA fingerprinting of parthenogenetic root-knot nematodes using AFLP analysis. *Molecular Ecology* 7: 119–125.
38. Blok VC (2005) Achievements in and future prospects for molecular diagnostics of plant-parasitic nematodes. *Canadian journal of plant pathology-revue canadienne de phytopathologie* 27: 176–185.
39. Agudelo P, Robbins RT, Stewart JMcD, Szalanski AL (2005) Intraspecific variability of *Rotylenchulus reniformis* from cotton-growing regions in the United States. *Journal of Nematology* 37: 105–114.
40. Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C(T)}$ method. *Methods* 25: 402–408.
41. Gao X, Hartman GL, Lambert KN, Li S (2004) Detection and quantification of *Fusarium solani* f. sp. *glycines* in soybean roots with real-time quantitative polymerase chain reaction. *Plant Disease* 88: 1372–1380.
42. Finetti-Sialer MM, Ciancio A (2005) Isolate-Specific Detection of Grapevine fanleaf virus from *Xiphinema* index through DNA-Based Molecular Probes. *Phytopathology* 95: 262–268.
43. Atallah ZK, Stevenson WR (2006) A methodology to detect and quantify five pathogens causing potato tuber decay using real-time quantitative polymerase chain reaction [electronic resource]. *Phytopathology* 96: 1037–1045.
44. Okubara PA, Schroeder KL, Paulitz TC (2008) Identification and Quantification of *Rhizoctonia solani* and *R. oryzae* Using Real-Time Polymerase Chain Reaction [electronic resource]. *Phytopathology* 98: 837–847.
45. Berry SD, Fargette M, Spaull VW, Morand S, Cadet P (2008) Detection and quantification of root-knot nematode (*Meloidogyne javanica*), lesion nematode (*Pratylenchus zaei*) and dagger nematode (*Xiphinema elongatum*) parasites of sugarcane using real-time PCR. *Molecular Cell Probes* 22: 168–176.
46. Showmaker K, Lawrence GW, Lu S, Balbalian C, Klink VP (2011) Quantitative field testing *Rotylenchulus reniformis* DNA from metagenomic samples isolated directly from soil. *PLoS ONE* 6 (12): e28954. doi:10.1371/journal.pone.0028954.
47. Klink VP, Martins VE, Overall CC, Alkharouf NW, MacDonald MH, et al. (2007) A decline in transcript abundance for *Heterodera glycines* homologs of *Caenorhabditis elegans* uncoordinated genes accompanies its sedentary parasitic phase. *BMC Developmental Biology* 7: 35.
48. van Megen H, van den Elsen S, Holterman M, Karssen G, Mooyman P, et al. (2009) A phylogenetic tree of nematodes based on about 1200 full-length small subunit ribosomal DNA sequences. *Nematology* 11: 927–950.
49. Alkharouf N, Klink VP, Matthews BF (2007) Identification of *Heterodera glycines* (soybean cyst nematode [SCN]) DNA sequences with high similarity to those of *Caenorhabditis elegans* having lethal mutant or RNAi phenotypes. *Experimental Parasitology* 115: 247–258.
50. Yik C-P, Birchfield W (1982) Reactions of sweet potato root tissue to the reniform nematode. *Plant Disease* 66: 707–709.
51. Daday E von (1905) Untersuchungen fiber die Siisserwasser-Mikrofauna Paraguays. *Zoologica, Stuttgart* 18: 1–349.
52. Bastian HC (1865) Monograph on the Anguillulidae or free nematodes, marine, land and freshwater, with description of 100 new species. *Transactions of the Linnaean Society of London*, 25: 73–184.
53. Zengel JM, Epstein HF (1980) Identification of genetic elements associated with muscle structure in the Nematode *Caenorhabditis elegans*. *Cell Motility* 1: 73–97.
54. Ritter ME (2009) *The Physical Environment: an Introduction to Physical Geography*.
55. Kofoid CA, White WA (1919) A new nematode infection of man. *Journal of the American Medical Association* 72: 567–569.
56. Chitwood BG (1949) Root-knot nematodes. I. A revision of the genus *Meloidogyne* Goeldi, 1887. *Proc. Helminthol. Soc. Wash.* 16: 90–104.
57. Matsye PD, Lawrence GW, Youssef RM, Kim K-H, Matthews BF, et al. (2012) The expression of a naturally occurring, truncated allele of an α -SNAP gene suppresses plant parasitic nematode infection. *Plant Molecular Biology* 80: 131–155.