

1-1-2008

## **Determination of Polyphenol Oxidase (PPO) Activity, Anthocyanin Contents and the Phytonutrient Changes in Blueberry Juice as Influenced by Different Processing Methods**

Jelena Stojanovic

Follow this and additional works at: <https://scholarsjunction.msstate.edu/td>

---

### **Recommended Citation**

Stojanovic, Jelena, "Determination of Polyphenol Oxidase (PPO) Activity, Anthocyanin Contents and the Phytonutrient Changes in Blueberry Juice as Influenced by Different Processing Methods" (2008). *Theses and Dissertations*. 1348.

<https://scholarsjunction.msstate.edu/td/1348>

This Dissertation is brought to you for free and open access by the Theses and Dissertations at Scholars Junction. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of Scholars Junction. For more information, please contact [scholcomm@msstate.libanswers.com](mailto:scholcomm@msstate.libanswers.com).

DETERMINATION OF POLYPHENOL OXIDASE (PPO) ACTIVITY,  
ANTHOCYANIN CONTENTS AND THE PHYTONUTRIENT  
CHANGES IN BLUEBERRY JUICE AS INFLUENCED BY  
DIFFERENT PROCESSING METHODS

By

Jelena Stojanovic

A Dissertation  
Submitted to the Faculty of  
Mississippi State University  
in Partial Fulfillment of the Requirements  
for the Degree of Doctor of Philosophy  
in Food Science and Technology  
in the Department of Food Science, Nutrition and Health Promotion

Mississippi State, Mississippi

August 2008

DETERMINATION OF POLYPHENOL OXIDASE (PPO) ACTIVITY,  
ANTHOCYANIN CONTENTS AND THE PHYTONUTRIENT  
CHANGES IN BLUEBERRY JUICE AS INFLUENCED BY  
DIFFERENT PROCESSING METHODS

By

Jelena Stojanovic

Approved:

---

Juan L. Silva  
Professor of Food Science, Nutrition and  
Health Promotion (Director of Dissertation  
and Graduate Coordinator)

---

Din-Pow Ma  
Professor of Biochemistry and  
Molecular Biology  
(Minor Professor and Minor  
Graduate Coordinator)

---

M. Wes Schilling  
Associate Professor of Food Science,  
Nutrition and Health Promotion  
(Committee Member)

---

Frank B. Matta  
Professor of Plant and Soil Sciences  
(Committee Member)

---

Charles H. White  
Professor Emeritus of Food Science,  
Nutrition and Health Promotion  
(Committee Member)

---

Melissa Mixon  
Interim Dean of the College of  
Agriculture and Life Sciences

Name: Jelena Stojanovic

Date of Degree: August 9, 2008

Institution: Mississippi State University

Major Field: Food Science and Technology

Major Professor: Dr. Juan L. Silva

Title of Study: DETERMINATION OF POLYPHENOL OXIDASE (PPO) ACTIVITY,  
ANTHOCYANIN CONTENTS AND THE PHYTONUTRIENT  
CHANGES IN BLUEBERRY JUICE AS INFLUENCED BY  
DIFFERENT PROCESSING TREATMENTS

Pages in Study: 177

Candidate for Degree of Doctor of Philosophy

Inhibition of blueberry PPO activity by sodium benzoate, potassium sorbate and potassium metabisulfite and their influence on degradation of individual anthocyanins in an extract was studied. Maceration of blueberries was carried out at 55°C for 1h with the addition of 0.1% sodium benzoate or with blanching pretreatment at 90°C for 1min. After maceration pretreatments the extracted juice was processed with traditional hot fill pasteurization, high hydrostatic pressure (HHP) and pulsed electric field (PEF).

Sodium benzoate and potassium metabisulfite were very effective PPO inhibitors in concentrations of 0.1% and 10ppm, respectively. Potassium sorbate was the weakest inhibitor, with 50% PPO remaining.

Degradation of anthocyanins by PPO was dependent on their structure. Tri-phenolic anthocyanins experienced the most degradation, followed by diphenolic and monophenolic compounds, respectively. Sodium benzoate was the most effective at

preventing anthocyanin degradation; potassium metabisulfite did not have any protective effect, while potassium sorbate increased anthocyanin degradation

Blanching of blueberries inactivated native PPO, but also increased the degradation of anthocyanins, especially malvidin glycosides. Addition of 0.1% sodium benzoate decreased PPO activity when compared to frozen blueberries but not in respect to control maceration.

Only 12% of anthocyanins and 33-41% of phenolics were extracted into juice from the frozen fruit. Hot fill pasteurization, high hydrostatic pressure and pulsed electric field did not significantly influence anthocyanins, phenolics and antioxidant activity in blueberry juice.

## DEDICATION

I would like to dedicate this research to my late cousin Natasa Vojnovic-Dakovic

## ACKNOWLEDGMENTS

I would like to acknowledge many people for helping me during my doctoral work. I would especially like to thank my advisor, Dr. Juan Silva, for his generous time and commitment. Throughout my doctoral work he encouraged me to develop independent thinking and helped me realize my true potentials.

I am also very grateful for having an exceptional doctoral committee and wish to thank Dr. Wes Schilling, Dr. Din-Pow Ma, Dr Frank Matta and Dr. Charles White for their continual support and encouragement.

I owe a special note of gratitude to Bill Holmes for setting up and teaching me how to use HPLC equipment and for helping me conduct analytical analysis at any time of the day.

Author would like to thank Dr. Aryana Kayanush for allowing me to use Pulsed Electric Field equipment at Louisiana State University and for his help in conducting PEF experiments. I am also very grateful to Dr. George Flick Jr. for conducting High Pressure experiments at Virginia Polytechnic Institute.

My sincere gratitude goes to Dr. Boris Stojanovic, “Professor”, and Dr. Elisabeth Stojanovic, “Ms Betty”, for being my family in Starkville. Their encouragement and support helped me through the toughest times.

I would also like to thank my family, my mother Cveta, my father Branko, my brother Radovan Stojanovic and my husband Aleksandar Todorovic for their patience and support.

I would like to acknowledge Ann and Roy Partridge for taking me into their home and being my American family.

I thank my friends Maria and Ovy Paraschivescu for their hospitality, encouragements and unselfish help any time when I need it.

I extend many thanks to my lab colleagues and friends, Dr. Kim, Lilly and Lu for making long days in the lab more pleasurable, and to the Department of Food Science staff, Donna Bland, Mary Andol and Joseph Andol.

I thank my long time friends Emina Kopic, Victorya Volkova and Yelena Syusko for their encouragement and friendship.

Last, but certainly not least, I would like to thank my friend Vesna Matijevic for making me happy when I really needed to smile and for offering her shoulder to cry on during tough times.

....

## TABLE OF CONTENTS

	Page
DEDICATION .....	ii
ACKNOWLEDGMENTS .....	iii
LIST OF TABLES .....	viii
LIST OF FIGURES .....	x
CHAPTER	
I. INTRODUCTION .....	1
II. LITERATURE REVIEW .....	9
Polyphenol oxidase .....	9
Blueberry PPO .....	14
Inhibition of PPO activity .....	15
Anthocyanins .....	23
Degradation of anthocyanins by PPO .....	26
Degradation of o-diphenolic anthocyanins .....	29
Degradation of non-diphenolic anthocyanins .....	30
Inhibition of anthocyanins degradation .....	32
Blueberry juice .....	35
Blueberry juice processing methods .....	37
PPO in food processing .....	39
Acids in blueberries .....	42
Phenolics .....	43
Changes in anthocyanins during juice processing .....	44
Changes in phenolics during juice processing .....	51
Non-thermal processing .....	53
Pulsed electric field .....	53
High pressure processing .....	57

III.	MATERIALS AND METHODS .....	61
	EXPERIMENT I .....	61
	Extraction of anthocyanins and phenolics .....	61
	Total anthocyanins .....	62
	Anthocyanin determination by HPLC-DAD and LC-MS .....	63
	Extraction of Polyphenol Oxidase (PPO) .....	64
	Determination of total protein.....	65
	PPO assay.....	66
	Inhibition study .....	67
	Degradation of anthocyanins by blueberry PPO in the model system.....	68
	Statistical analysis.....	69
	EXPERIMENT II .....	70
	Maceration pretreatment of blueberry mash .....	70
	Blueberry juice filtration.....	70
	Juice treatments.....	71
	Statistical analysis.....	74
	Analysis.....	74
	Extraction of anthocyanins and phenolics .....	74
	Anthocyanin determination by HPLC-DAD .....	74
	Extraction and assay of polyphenol oxidase (PPO).....	75
	Total phenolics.....	75
	Polymeric color.....	76
	Antioxidant activity .....	77
	Acids .....	78
IV.	RESULTS AND DISCUSSION .....	81
	EXPERIMENT I .....	81
	Enzyme activity and inhibition .....	81
	Identification of anthocyanins .....	84
	Degradation of anthocyanins.....	86
	EXPERIMENT II.....	116
	Identification and quantification of anthocyanins.....	114
	Identification and quantification of acids .....	115
	PPO activity in blueberry mash .....	116
	Degradation of anthocyanins, acids and phenolics in blueberry mash ....	117
	Juice yield .....	121
	Anthocyanins, phenolics and acids in pressed juice and press cake.....	121
	Juice processing .....	126
	Pulsed Electric Field, PEF .....	126
	Pasteurization.....	128
	High pressure processing, HPP.....	129

V. SUMMARY AND CONCLUSIONS .....	159
REFERENCES .....	165

## LIST OF TABLES

TABLE	Page
1. Linear gradient program used for the separation of anthocyanins .....	63
2. Concentration of inhibitors used in the PPO inhibition study .....	68
3. Slopes of the inhibitor regression lines .....	92
4. Retention times, % area and m/z of total and aglucone of identified Anthocyanins in blueberry extract using High Pressure Liquid Chromatography .....	95
5. Concentration of individual anthocyanins expressed as cyanidin 3-glucoside as affected by different maceration treatments .....	134
6. Concentration of acids in blueberry mash as affected by different maceration treatment .....	137
7. Concentration of individual anthocyanins expressed as cyanidin 3-glucoside in pressed juice as affected by different maceration treatments .....	140
8. Concentration of acids in pressed juice as affected by different maceration treatment .....	143
9. Concentration of individual anthocyanins expressed as cyanidin 3-glucoside in pomace as affected by different maceration treatments .....	146
10. Concentration of acids in blueberry pomace as affected by different treatments .....	148
11. Concentration of individual anthocyanins, total phenolic, % polymeric color, total anthocyanins and antioxidant activity before and after PEF treatment .....	150

12.	Concentration of individual anthocyanins, total phenolic, % polymeric color, total anthocyanins and antioxidant activity before and after pasteurization .....	153
13.	Concentration of individual anthocyanins expressed as cyanidin 3-glucoside in juice as affected by different processing pressures .....	156
14.	Concentration of acids in juice as affected by high pressure treatments.....	157

## LIST OF FIGURES

FIGURE	Page
1. Structure of benzoic acid .....	16
2. Structure of cinnamic acid.....	16
3. Structure of most commonly found anthocyanidins.....	24
4. Forms of anthocyanins present in acidic solution .....	25
5. Standard curve for determination of total protein with BSA as a standard...	65
6. Digital controller model 10 (a) and Rank inverted oxygen electrode (b).....	67
7. Schematic representation of blueberry mash maceration treatments .....	72
8. OSU-4K Pulsed electric field equipment .....	73
9. Cyanidin 3-glucoside standard curve .....	75
10. Gallic acid standard curve .....	76
11. 2,2 Diphenyl-1-picrylhydrazyl (DPPH) standard curve.....	78
12. Quinic acid standard curve .....	79
13. Malic acid standard curve.....	79
14. Citric acid standard curve.....	80
15. Shikimic acid standard curve.....	80
16. Effect of sodium benzoate and potassium sorbate on blueberry PPO activity.....	92
17. Effect of potassium metabisulfite on PPO blueberry activity .....	93

18.	Effect of combination of sodium benzoate and potassium metabisulfite on blueberry PPO activity .....	93
19.	HPLC chromatogram of 14 anthocyanins detected at 520 nm using DAD ..	94
20.	Extracted ion chromatogram of 15 identified anthocyanins.....	94
21.	Extracted chromatogram of delphinidin 3-galactoside and delphinidin 3-glucoside m/z 465 .....	96
22.	Ion chromatogram of delphinidin 3- galactoside (peak #1) .....	96
23.	Ion chromatogram of delphinidin 3- glucoside (peak #2) .....	96
24.	Extracted chromatogram of cyanidin 3-galactoside, cyanidin 3- glucoside and petunidin 3-arabinoside m/z 449 .....	97
25.	Ion chromatogram of cyanidin 3- galactoside (peak #3).....	97
26.	Ion chromatogram of cyanidin 3- glucoside (peak #5) .....	97
27.	Ion chromatogram of petunidin 3- arabinoside (peak #10).....	98
28.	Extracted chromatogram of delphinidin 3-arabinoside m/z 435 .....	98
29.	Ion chromatogram of delphinidin 3- arabinoside (peak #4).....	98
30.	Extracted chromatogram of petunidin 3-galactoside and petunidin 3- glucoside m/z 479 .....	99
31.	Ion chromatogram of petunidin 3- galactoside (peak #6) .....	99
32.	Ion chromatogram of petunidin 3- glucoside (peak #8) .....	99
33.	Extracted chromatograms of peonidin 3-galactoside, peonidin 3- glucoside and malvidin 3-arabinoside m/z 463 .....	100
34.	Ion chromatogram of peonidin 3- galactoside (peak #9).....	100
35.	Ion chromatogram of peonidin 3- glucoside (peak #11) .....	100
36.	Ion chromatogram of malvidin 3- arabinoside (peak #14).....	101

37.	Extracted chromatograms of malvidin 3-galactoside and malvidin 3- glucoside m/z 493 .....	101
38.	Ion chromatogram of malvidin 3-galactoside (peak #12) .....	101
39.	Ion chromatogram of malvidin 3-glucoside (peak #13).....	102
40.	Extracted chromatogram of peonidin 3-arabinoside m/z 433 .....	102
41.	Ion chromatogram of peonidin 3-arabinoside .....	102
42.	Degradation of total anthocyanins by blueberry PPO with and without inhibitors .....	103
43.	Degradation of anthocyanins by blueberry PPO .....	103
44.	Degradation of anthocyanins by PPO grouped according to the sugar in the molecule .....	104
45.	Degradation of delphinidin anthocyanins by PPO with and without inhibitors .....	104
46.	Degradation of delphinidin 3-galactoside by PPO with and without inhibitors .....	105
47.	Degradation of delphinidin 3-glucoside by PPO with and without inhibitors .....	105
48.	Degradation of delphinidin 3-arabinoside by PPO with and without inhibitors .....	106
49.	Degradation of petunidin anthocyanin by PPO with and without inhibitors .....	106
50.	Degradation of petunidin 3-galactoside by PPO with and without inhibitors .....	107
51.	Degradation of petunidin 3-glucoside by PPO with and without inhibitors .	107
52.	Degradation of petunidin 3-arabinoside by PPO with and without inhibitors .....	108
53.	Degradation of cyanidin anthocyanins by PPO with and without inhibitors .....	108

54.	Degradation of cyanidin 3-galactoside by PPO with and without inhibitors .....	109
55.	Degradation of cyanidin 3-glucoside by PPO with and without inhibitors...	109
56.	Degradation of cyanidin 3-arabinoside by PPO with and without inhibitors .....	110
57.	Degradation of malvidin anthocyanins by PPO with and without inhibitors .....	110
58.	Degradation of malvidin 3-galactoside by PPO with and without inhibitors .....	111
59.	Degradation of malvidin 3-glucoside by PPO with and without inhibitors ..	111
60.	Degradation of malvidin 3-arabinoside by PPO with and without inhibitors .....	112
61.	Degradation of peonidin anthocyanins by PPO with and without inhibitors .....	112
62.	Degradation of peonidin 3-galactoside by PPO with and without inhibitors .....	113
63.	Degradation of peonidin 3-glucoside by PPO with and without inhibitors...	113
64.	HPLC separation of acid standards detected at 214 nm .....	131
65.	HPLC separation of acids detected at 214 nm.....	131
66.	Concentration of total anthocyanins after 60 min of maceration as affected by different treatments .....	132
67.	Concentration of anthocyanins aglucones after 60 min of maceration as affected by different treatments .....	133
68.	Concentration of total phenolics after 60 min of maceration as affected by different treatments .....	135
69.	Changes in polymeric color after 60 min of maceration as affected by different treatments .....	136
70.	Activity of PPO ( $\mu\text{mol O}_2/\text{min}/100\text{g}$ ) after 60 min of maceration as affected by different treatments .....	137

71.	Concentration of total anthocyanins in the pressed juice as affected by different treatments .....	138
72.	Concentration of anthocyanin aglucones in the pressed juice as affected by different treatments .....	139
73.	Concentration of total phenolic in the pressed juice expressed as gallic acid equivalents as affected by different maceration treatments .....	141
74.	Percent polymeric color in the pressed juice as affected by different maceration treatments .....	142
75.	Yield of the pressed juice as affected by different maceration treatments....	143
76.	Concentration of total anthocyanins in pomace as affected by different maceration treatments .....	144
77.	Concentration of anthocyanins aglucones in pomace as affected by different maceration treatments .....	145
78.	Concentration of total phenolics in pomace expressed as gallic acid equivalents as affected by different maceration treatments .....	147
79.	Percent polymeric color in pomace as affected by different maceration treatments .....	148
80.	Concentration of total anthocyanins before and after PEF treatment.....	149
81.	Spectrophotometric recordings of the disappearance of DPPH at 515 nm in the presence of blueberry juice before and after PEF treatment.....	151
82.	Concentration of acids before and after PEF treatment.....	151
83.	Concentration of anthocyanin aglucones before and after pasteurization ....	152
84.	Spectrophotometric recordings of the disappearance of DPPH at 515 nm in the presence of blueberry juice before and after pasteurization .....	152
85.	Concentration of acids before and after pasteurization .....	154
86.	Concentration of total anthocyanins as affected by different processing pressures.....	154

87.	Concentration of anthocyanin aglucones as affected by different processing pressures.....	155
88.	Percent anthocyanin polymerization as affected by different processing pressures.....	155
89.	Effect of different processing pressures on total phenolics expressed as ppm of gallic acid equivalents .....	157
90.	Disappearance of DPPH, at 515 nm, in blueberry juice processed at different pressures for 10 min .....	158

## CHAPTER I

### INTRODUCTION

Polyphenol oxidase (PPO) is a generic term for a group of enzymes that catalyzes the oxidation of phenolics compounds leading to the development of browning (Kader and others 1997b). The extent to which naturally occurring phenolic substrates contribute to enzymatic browning of individual fruits and vegetables depends on the localization and concentration of the phenols as well as on the color intensity of the macromolecular pigments obtained from the different quinines (Vamos-Vigyazo 1981).

O-quinones, the primary products of the oxidative reaction catalyzed by the enzyme, (a) react with each other to form high molecular weight polymers, (b) form molecular complexes with amino acids or proteins, and anthocyanins, and (c) oxidize compounds that have lower oxidation-reduction potential (Mathew and Parpia 1971). These subsequent reactions may bring about changes in physical, chemical, nutritional and sensory characteristics of food products (Mayer and Harel 1979).

Browning may be prevented not only by inactivating the enzyme, but also by eliminating one of the two substrates necessary for the reaction (Phenols and  $O_2$ ), or by reacting with the products of enzyme action to inhibit the formation of the colored compounds produced in secondary, non-enzymatic reaction steps (Vamos-Vigyazo 1981).

Many compounds have been identified as possible PPO and browning inhibitors but most of them are not allowed for the use in food. The use of browning inhibitors in food processing is restricted by special requirements such as non-toxicity, wholesomeness, effect on taste, flavor, texture, etc (Vamos-Vigyazo 1981).

Carboxylic acids are shown to be strong inhibitors of PPO (Pifferi and others 1974; Walker and Wilson 1975; Walker and McCallion 1980; Gunata and others 1987; Ferrar and Walker 1996). Benzoic acid and some derivatives of cinnamic acid were found to be mostly competitive inhibitors of PPO (Pifferi and others 1974).

Sulfites are arguably the most versatile food additives available. They act not only as a food preservative but have an important role as inhibitors of enzymatic and non-enzymatic browning as inhibitors of a wide range of enzymes including proteases, oxidases, peroxidases, and are alleged to prevent oxidative spoilage (Wedzicha 1992). Sulfites are highly effective in controlling browning but are subject to regulatory restrictions because of adverse effect on health (Sapers 1993). Sulfites can act as PPO inhibitors and can also react with intermediates to prevent pigment formation (Sayavedra-Soto and Montgomery 1986)

Anthocyanins are the most important water-soluble pigments in plant tissues, and produce blue, red and purple colors. Color of anthocyanins-containing media depends on structure and concentration of the pigment, pH, temperature, presence of copigments, metallic ions, enzymes, oxygen, ascorbic acid, sugars and their degradation products, sulfur dioxide, actual anthocyanins concentration and other factors (Mazza and Miniati 1993). There is experimental evidence that certain anthocyanins and flavonoids have anti-inflammatory properties, and there are reports

that orally administered anthocyanins are beneficial for treating diabetes, ulcers and may have antiviral and antimicrobial activities and can also provide protection against UV radiation (Mazza and Miniati 1993). Anthocyanins, as a functional food component, can aid in the prevention of obesity and diabetes (Tsuda and others 2003). It is well known that anthocyanins are poor substrates of PPO, and this is probably due to the presence of the sugar moiety causing steric hindrance, since the aglucon forms are often oxidized by PPO (Mathew and Parpia 1971).

Blueberries are a very rich source of anthocyanins and other phenolics. Enzymatic oxidation of phenolic compounds present in blueberries by PPO can occur during processing as soon as the berries are damaged. The first reaction step that occurs is oxidation of phenolic substrates to their o-quinones. Quinones are very reactive species which are able, as oxidants, to oxidize other substrates, being reduced in the process to the original phenol, and as electrophiles to react with various nucleophiles (Sarni and others 1995). These reactions of o-quinones, namely oxidation and condensation reactions can lead to discoloration and can involve other phenolic compounds especially anthocyanins. The use of model systems has shown that the enzymatically generated o-quinones play an essential role in the process of anthocyanin degradation (Kader and others 1999a). Degradation of anthocyanins can be inhibited by direct inhibition of PPO and with that formation of o-quinones or by the inhibiting anthocyanins from reacting with already generated o-quinones. Maintaining high antioxidant activity with preserving anthocyanins and other phenolics is of crucial importance for any blueberry processed product. Inhibition of PPO during processing, thus preventing oxidation of phenolics and degradation of

anthocyanins should be a preliminary step in any blueberry processing. Since anthocyanins are heat sensitive use of inhibitors could be the best solution for PPO inhibition without the degradation of anthocyanins.

Health attributes ascribed to blueberries create a need for the development of processed blueberry products that are convenient for the consumer. Blueberries can be pressed into juice and juice concentrate, in order to reach a more wide spread consumption (Rossi and others 2003). Desirable color is a very important sensory characteristic for the consumer of fruits and processed fruit products (Garzon and Wrolstad 2002). Huge economic losses can also be accrued by the company due to the loss of attractive product color. A concord grape juice producer reported about \$ 700,000 in losses due to the loss of juice color (Wrolstad and others 1994). The stability of color in blueberries is an important aspect of quality control during processing and storage (Yang and Yang 1987). Not just because they impart color but also because of their possible beneficial effect, particular attention has to be paid to the changes that anthocyanin pigments undergo during processing.

Anthocyanins are not as efficiently extracted in the pressing operation as sugars, acids and other water solubles, which can have a negative impact on juice quality (Skrede and others 2000; Lee and others 2002). The anthocyanins that give color to blueberries exist almost exclusively in the skin, whereas phenolics and other antioxidants are mostly in the flesh (Lee and Wrolstad 2004). Since anthocyanins in blueberries are only found in the skin, skin breaking and maceration are needed to extract color and obtain good quality juice. Lee and others (2002) concluded that heating of berries may have contributed to a breakdown of the skins, increasing

extraction of the color, but still with a substantial amount of anthocyanins left in the press cake. Cold processing of blueberries was not as efficient at transferring antioxidants or phenols to the final juice as hot processing (Carlson 2003).

During juice processing, anthocyanins can be lost due to enzymatic breakdown, heat treatment and some are removed with the pulp (Iversen 1999). The primary steps of juice processing (thawing, crushing, depectinization and pressing) contribute to a large loss in total anthocyanins (Skrede and others 2000; Lee and others 2002). The studies listed above reported that substantial losses of anthocyanins and other polyphenolics occurred when blueberries were processed into juice, and in addition different classes of compounds had varying susceptibility to degradation with different processing operations, and the highest losses occurred during milling and depectinization due to the action of native polyphenoloxidase, PPO (Skrede and others 2000).

Processing or any wounding of fruits and vegetables may cause cell disruption that can lead to quinone formation due to the action of PPO leading to product deterioration. Not only the appearance of food and beverages may be affected, but also their taste and nutritional value, often decreasing the quality of the final products (Gandia-Herrero and others 2005). Degradation of anthocyanins occurs when active PPO and its substrate are present in the system. Although anthocyanins are not direct substrates for PPO, anthocyanins with a *o*-diphenolic B ring were oxidized via enzymatically generated *o*-quinines of catechol, catechin, chlorogenic acid (Peng and Markakis 1963; Pifferi and Cultrera 1974; Raynal and Moutounet 1989; Wesche-Ebeling and Montgomery 1990b) and caftaric acid (Cheynier and others 1994). It has

been reported that heat inactivation of PPO is unacceptable in anthocyanin-containing juice products, since the high temperature required for PPO inactivation can cause anthocyanin degradation (Siddiq and others 1992).

Processing and storage can also have marked effects on the phenolics' content of fruits that might also affect their health promoting properties (Zafrilla and others 2001). As a general rule, during processing, phenolic biosynthesis is interrupted by the enzyme's destruction and/or cell structure degradation (Tomas-Barberan and Espin 2001a). Processing can enhance phenolic compound degradation (chemically or enzymatically if the oxidative enzymes have not been inactivated) or can produce chemical changes that affect quality characteristics (Tomas-Barberan and Espin 2001a).

Blanching or addition of PPO inhibitors can have a positive effect on juice quality. Blanching of strawberries before juice and concentrate production had a protective effect on anthocyanin pigments, leucoanthocyanins, flavanols, total phenolics and ascorbic acid, and also resulted in improved color stability (Wrolstad and others 1980). Chemical treatments like SO<sub>2</sub>, citric acid and SnCl<sub>2</sub> had a stabilizing effect on strawberry anthocyanins during processing of strawberry jam (Sistrunk and others 1982). Heat and SO<sub>2</sub> pretreatment before pressing the juice increased recovery of anthocyanins in all processing steps (Lee and others 2002).

There is a demand from consumers and retailers for minimum processed foods that are also safe. Therefore, producers and researchers are searching for alternative food processing methods to gently preserve foods (Houska and others 2006).

Although thermal preservation methods provide safe foods, there is loss of food quality, and nutritional value that is associated with this processing method. Hence, the main objective of low- and non-thermal preservation methods is to minimize the degradation of food quality through limiting heat damage (Senorans and others 2003). Compared with thermal processing, nonthermal methods provide a high retention of flavors and nutrients, giving products a “natural/fresh” taste (Zhong and others 2005).

High voltage pulsed electric fields (PEF) is a promising non thermal processing technology. Treating liquid foods with PEF may inactivate microorganisms and enzymes with only small increases in temperature, simultaneously providing consumers with safe, nutritious, and fresh foods (Aguilar-Rosas and others 2007). Consumer demand for a higher quality, fresh appearing and safe food supply is the ultimate catalyst for the emergence of PEF on a commercial scale (Yeom and others 2002)

Cranberry juice treated by PEF had a similar flavor or aroma profile as the nontreated juice; whereas thermal treatment significantly altered the overall flavor profile of the juice (Jin and Zhang 1999). No differences were observed in the content of anthocyanin pigments between PEF-treated samples and controls. However, thermal treatment significantly reduced their anthocyanin content

High hydrostatic pressure (HHP) is considered a nonthermal technology with the most promising perspective of industrial utilization. There are already known industrial applications in Japan, USA, France and Spain (Houska and others 2006). This technology can enable processors to produce innovative foods with fresh-like, natural-like attributes and natural looking colors which are all aspects that are valued

by consumers (Deliza 2005). In addition information on the benefits that are offered by high pressure technology presented on the juice package had a positive contribution on purchase intention (Deliza 2005). High pressure treatment is expected to be less detrimental than thermal processing to low molecular weight food compounds such as flavoring agents, pigments, vitamins, etc., since covalent bonds are not affected by pressure ( Butz and others 2002). The pressure stability of antioxidants is of interest since they play an important role in reducing the risk of free radical-related oxidative damage associated with a number of diseases (Butz and others 2002).

High hydrostatic pressure is a promising alternative to traditional thermal processing techniques of food preservation, but associated changes to a diversity of phytonutrients have not been extensively investigated (Talcott 2003). Making comparisons between HHP and thermal processes is one way to assess its benefits by monitoring destruction of heat labile compounds (Talcott 2003).

The need to preserve valuable phytonutrients in blueberries with minimum impact on the quality and taste in the final product leads to the need for this type of study. The objectives of this study were:

1. To determine inhibition of polyphenol oxidase (PPO) by preservatives and their influence on anthocyanins in a blueberry extract and
2. To determine influence of blueberry mash pretreatment and various processes on anthocyanins, phenolics and antioxidant activity of blueberry juice.

## CHAPTER II

### LITERATURE REVIEW

#### **Polyphenol oxidase**

Polyphenol oxidase (PPO) is a generic term for the group of enzymes that catalyze the oxidation of phenolic compounds that cause browning. Based on substrate specificity there are: Monophenol monooxygenase, cresolase or tyrosinase (EC 1.14.18.1); Diphenol oxidase or catechol oxidase (EC 1.10.3.2); and laccase or p-diphenol oxygen oxidoreductase (EC 1.10.3.1) (Shahidi and Naczki 1995). Numerous reports on enzymes from various sources suggests that there is a copper content of one atom per polypeptide chain or subunit of catechol oxidase (Mayer and Harel 1979). The structure of the active site of the enzyme, in which copper is bound by six or seven histidine residues and a single cysteine residue is highly conserved (Mayer 2006).


A variety of phenolic compounds can serve as PPO substrates. The compartmentation of the phenolic substrate of the enzyme, both in special cells and within the cell have been reported which results in the separation between the enzyme and the bulk of its phenolic substrates *in situ* (Mayer and Harel 1979).

The most important natural substrates of PPO in fruits and vegetables are catechins, cinnamic acid esters (chlorogenic acid), 3,4-dihydroxy phenylalanine


(DOPA), and tyrosine (Vamos-Vigyazo 1981). The sphere of naturally occurring substrates of PPO is limited by the fact that the enzyme does not act on glycosides (Baruah and Swain 1959), and most of the phenolic in plants are present in that form. Negative or positive charges near the phenolic ring, disturb the catalytic process resulting in low activity (Casado-Vela and others 2005). Affinity of the PPO toward substrate is highly dependent on the plant source from which the enzyme is extracted. The crude enzyme extract from concord grapes had activity toward all of the dihydroxyphenols but no activity with monophenols (Cash and others 1976). Although gallic and chlorogenic acid were of dihydroxy configuration, the crude grape enzyme extract did not effectively utilize either of these compounds as its substrate (Cash and others 1976) while the chlorogenic acid was the best substrate for blueberry PPO (Kader and others 1997b). The extent to which naturally occurring phenolic substrates contribute to the enzymatic browning of individual fruits and vegetables depends on the localization and concentration of the phenols as well as on the color intensity of the macromolecular pigments that are obtained from the different quinines (Vamos-Vigyazo 1981).

Polyphenol oxidase catalyzes two basic reactions (Shahidi and Naczk 1995):

1. hydroxylation to the *o*-position adjacent to an existing hydroxyl group of the phenolic substrate (monophenol oxidase activity).



2. and oxidation of diphenol to *o*-benzoquinones (diphenol oxidase activity).



Both reactions utilize molecular oxygen as a co-substrate. Whether a single enzyme system exhibits both mono- and di phenol oxidase activities is still unclear. However, when both monophenol and diphenol oxidases are present in plants, the ratio of monophenol to diphenol oxidase activity is usually 1:10 or as low as 1:40 (Nicolas and Potus 1994).

The primary products of the oxidative reaction catalyzed by the enzyme, the *o*-quinones, (a) react with each other to form high molecular weight polymers, (b) form molecular complexes with amino acids or proteins, and anthocyanins, and (c) oxidize compounds of lower oxidation-reduction potential (Mathew and Parpia 1971). Non-enzymatic reactions (a) and (b) lead to the formation of brown pigments, the

color which is darker the higher their molecular mass (Vamos-Vigyazo 1981). These subsequent reactions may bring about changes in physical, chemical, nutritional and sensory characteristics of food products (Mayer and Harel 1979). The rate of enzymatic browning reactions depends on the nature and content of phenolic compounds, activity of phenoloxidase present in food, presence of oxygen reducing substances and metal ions, pH and temperature (Shahidi and Naczk 1995).

Products of reaction of type (c) are colorless (Vamos-Vigyazo 1981). In the reaction that belong to group (c), quinines formed by the reaction of PPO, oxidize compounds of lower oxidation-reduction potentials and are again reduced to dihydroxyphenols which provides “fresh” substrate for PPO. This reaction can continue until the enzyme gets inactivated by the reaction products, or compounds of lower oxidation-reduction potential are depleted (Vamos-Vigyazo 1981).

PPO can be found in both soluble and membrane-bound forms in chloroplast, mitochondria, microsomes, peroxisomes and cytoplasm (Tomas-Barberan and Espin 2001b). Serradell and others (2000) concluded that strawberry PPO is associated with the membranes since it was released from the fruit using detergent (Triton X-100) and NaCl. The strength of binding of PPO to the membrane appears to be dependent on the tissue and the stage of development (Mayer and Harel 1979). Solubilization of PPO occurs during ripening or aging of fruits (Mayer and Harel 1979). It was strongly suggested that apple PPO is solubilized and denatured at the later stage of development, but with enough remaining to cause browning (Murata and others 1995). Serradell and others (2000) reported that PPO is more abundant in immature strawberry fruit than in the ripe one.

One of the oldest suggestions of the physiological role of PPO is that of o-diphenol synthesis. This suggestion is based on the ability of many catechol-oxidase preparations to oxidase monophenols to the corresponding o-quinones (Mayer and Harel 1979). Changes in the catecholase activity during maturation are related to the formation of anthocyanins (Sanchez-Ferrer and others 1989). PPO has also been reported to be a defensive protein against pathogen attack in various crops (Tomas-Barberan and Espin 2001b).

As with any other enzyme activity, PPO is highly dependent on pH and temperature. The optimum pH of PPO activity varies with the source of the enzyme and with the substrate in a relatively wide range but in most cases is between 4.0 and 7.0 (Vamos-Vigyazo 1981). Optimum activity of the crude enzyme extract occurred at pH 5.9-6.3 while over 50% of its maximum activity was retained at 3.4, the normal pH of the grape juice (Cash and others 1976). Polyphenol oxidase from strawberries showed maximum activity at a pH of 5.3 (Serradell and others 2000).

PPO is not an extremely heat-stable enzyme. Short exposures, in the tissue and solutions, to temperatures of 70 to 90°C are, in most cases, sufficient for partial or total irreversible destruction of its catalytic function (Vamos-Vigyazo 1981). Most PPO enzymes show optimum activity at temperatures between 30 and 40°C (Lamikanra and others 1992). Strawberry polyphenol oxidase exhibited maximum activity at 50°C and then decreased sharply at higher temperatures although it showed considerable activity (15-20%) at extreme temperatures of 0 to 70°C. These authors also reported that the enzyme retained 71 and 43% of its activity after incubation at 55°C for 30 and 120 min respectively, while temperatures of 65°C caused an almost

complete loss of activity even after 30 min. PPO from concord grapes showed a maximum activity between 25 and 30°C, after which the reaction rate declined very rapidly as temperature increased (Cash and others 1976). Enzyme from Emir grape extract was heat stable at 65°C, retaining up to 73% of activity after 30 min of heating with a D value of 213 min (Unal and Sener 2006). Activity of apple PPO increased with mild heating and reached a maximum, and as heating progressed, activity decreased, first gradually and then rapidly (Yemenicioglu and others 1997). This increased activity by heat was explained by the activation of latent PPO. The activation of latent PPO can be attributed to protein association and dissociation (Mathew and Parpia 1971).

### **Blueberry PPO**

Kader and others (1997b) used native PAGE to demonstrate that blueberry PPO presents two isoenzymes PPO1 and PPO2. Although they found that caffeic acid as a substrate would provide highest PPO activity, absence of caffeic acid in highbush blueberries would make chlorogenic acid a better substrate for assessing PPO activity in blueberries since it is the major hydroxycinnamic derivative. Oxidation of chlorogenic acid by purified blueberry PPO produced degradation of only 50%, whereas 70% oxidation was obtained when nonpurified PPO was used (Kader and others 1998). These researchers concluded that PPO in the nonpurified extract is probably protected by non-PPO proteins which react with the oxidized polyphenols, thereby sparing the PPO.

The optimum activity of the enzyme was a pH of 4 with the shoulder at pH 5 (Kader and others 1997b). At pH 5, about 80% of the maximum activity remained with the rapid decrease in activity with an increase in pH to 5.5 (Kader and others 1997b). This behavior was explained by the presence of two isoforms. Blueberry PPO was found to be very active at low pH's that are similar to the pH of fruits (3-3.2) (Kader and others 1994).

### **Inhibition of PPO activity**

Browning may be prevented by inactivating the enzyme, by eliminating one of the two substrates necessary for the reaction (Phenols and O<sub>2</sub>), or by reacting with the products of enzyme action to inhibit the formation of the colored compounds that are produced in secondary, non-enzymatic reaction steps (Vamos-Vigyazo 1981). In many cases, it is not easy to distinguish the different mechanisms that underlay the action of a browning inhibitor; moreover, some inhibitors act simultaneously on the enzyme and the substrate or products (Vamos-Vigyazo 1981). In general PPO inhibitors can be divided into two groups; those that act primarily on the enzyme and the those that react with the reaction products on the substrate. The use of browning inhibitors in food processing is restricted by special requirements such as non-toxicity, wholesomeness, effect on taste, flavor, texture, etc (Vamos-Vigyazo 1981).

Carboxylic acids are strong inhibitors of PPO (Pifferi and others 1974; Walker and Wilson 1975; Walker and McCallion 1980; Gunata and others 1987; Ferrar and Walker 1996). Benzoic acid (Figure 1) and some derivatives of cinnamic acid (Figure 2) are mostly competitive inhibitors of PPO (Pifferi and others 1974). The

only efficient structural requirement for the inhibitory effect is the association of the carboxyl group with the benzene nucleus or with an unsaturated open chain (Pifferi and others 1974). It was shown that PPO was inhibited by cinnamic and p-coumaric acids (Walker and Wilson 1975; Walker and McCallion 1980; Gunata and others 1987), benzoic acid (Gunata and others 1987) and to a lesser extent, by ferulic acid (Walker and Wilson 1975; Walker and McCallion 1980).




Figure 1. Structure of benzoic acid




Figure 2. Structure of cinnamic acid.

Hydroxylation, methylation of the benzene ring of these acids and the esterification of the carboxyl group of benzoic and cinnamic acids leads to a considerable decrease in their inhibitory strength (Gunata and others 1987). It has been suggested that for strong inhibition, aromatic acid inhibitors require a free carboxylic group to be substituted directly on the benzene ring (Vamos-Vigyazo 1981). Benzoic acid was found to be a competitive inhibitor when 4-methylcatechol and (+)-catechin were used as substrates and non-competitive with caffeic acid (Gunata and others 1987). Janovitz-Klapp and others (1990) also concluded that benzoic acid is competitive with 4-methyl catechol since it does not affect the apparent  $V_m$  and increases the apparent  $K_m$ . In the system with benzoic acid, phenolic substrate (4-methyl catechol) and oxygen are present, oxygen is the first substrate to be bound to apple PPO, forming an E-O<sub>2</sub> complex to which the benzoic acid as an inhibitor can bind only with enzymatic forms of PPO which are free of other phenolics i.e., E or E-O<sub>2</sub> (Janovitz-Klapp and others 1990b). Some PPO inhibitors require protonation of the acidic group (i.e., low pH) in order to display inhibition (Pifferi and others 1974; Janovitz-Klapp and others 1990a) that should be taken into account when investigating inhibition (Ferrar and Walker 1996). The inhibition of mushroom PPO by benzoic acid and by cyanide, respectively, showed the former to be competitive with catechol (substrate) and non competitive with O<sub>2</sub>, while the later inhibitor showed a reversed behavior. This was interpreted as the result of the existence of two distinct substrate binding sites on the enzyme molecule, one of which had a high affinity for aromatic compounds, including phenolic substrates, while the other, which probably contained the enzyme copper, served for

metal-binding agents and oxygen (Duckworth and Coleman 1970). Benzoic acid was found to be a mixed-type inhibitor of gum arabic PPO showing the strongest inhibition among all carboxylic acids tested at a pH of 5.3 (Billaud and others 1996).

The inhibitors of enzymatic browning that react with the reaction products or substrates, can be divided into following groups (Vamos-Vigyazo 1981).

1. Reducing agents acting on the formed quinines by restituting the o-dihydroxy phenols. These compounds are consumed in the process of inhibition, and thus provide only temporary protection against discoloration, unless used in high concentration, in which case reaction inactivation of the enzyme might occur prior to the depletion of reducing agent. Some frequently used representatives of this group are ascorbic acid, SO<sub>2</sub>, potassium metabisulfite, 2-mercaptoethanol, and thioglycollate.
2. Quinone couplers forming stable colorless compounds with quinones, thus providing permanent protection as long as they are not entirely consumed. Cysteine, glutathione, benzenesulphinic acid, DIECA, Na-ethyl xanthate , among others, are able to perform such reactions.

Sulfites are arguably the most versatile food additives available. They act not only as a food preservative but have an important role as inhibitors of enzymatic and non-enzymatic browning, inhibitors of a wide range of enzymes including proteases, oxidases, peroxidases, and are alleged to prevent oxidative spoilage (Wedzicha 1992). Sulfites are highly effective at controlling browning but are subject to regulatory restrictions because of adverse effect on health (Sapers 1993). The main reason for

the reactivity of sulfites in foods is the high nucleophilicity of the sulfite ion (Wedzicha 1992)

Sulfites have a very complex action on the PPO-polyphenol system (Vamos-Vigyazo 1981). Sulfites can act as PPO inhibitors and can also react with intermediates to prevent pigment formation (Sayavedra-Soto and Montgomery 1986). Sulfites can inhibit browning by combining irreversibly with the quinones to form colorless addition products, at the same time reducing the activity of the enzyme towards both mono- and dihydroxy phenols (Embs and Markakis 1965 and Markakis and Embs 1966). Inhibition of browning involves nucleophilic attack by the sulfite ion on formed quinone by PPO reaction to form sulfonate, where quinone is reduced and sulfonate is unreactive towards PPO (Wedzicha and others 1991). Inhibition of browning is permanent if all the substrate (quinone) is converted to sulfonate (Wedzicha and others 1991). Some authors have reported that sulfite has direct effect on enzyme (Golan-Goldhirsh and Whitaker 1984; Sayavedra-Soto and Montgomery 1986; Valero and others 1992). The product (quinone) accumulation curve in the presence of metabisulfite includes initial an lag period due to the chemical reaction between the quinone products of the enzyme catalysis and metabisulfite and a steady state phase, where all of the inhibitor has been depleted. During the steady state phase the catalytic activity that is expressed is lower than in the absence of metabisulfite, which is probably an indication of the direct effect of metabisulfite on PPO (Valero and others 1992). It is assumed that the inactivation of the oxy form of PPO involves the formation of an enzyme-sulfite complex, which subsequently undergoes an irreversible isomerization reaction that inactivates the enzyme (Valero and others

1992). The main action of sulfite on the enzyme might be sulfitolysis at the vital point for the enzyme activity (Sayavedra-Soto and Montgomery 1986). These researchers also suggested that the formation of PPO-SO<sub>3</sub> complex could have occurred due to the interaction between sulfite and PPO, forming inactive PPO that differed in some properties that was shown by additional bands on electrophoresis and the disappearance of the protein band where the active PPO should have been noticed (Sayavedra-Soto and Montgomery 1986). Presence of the lag phase was also observed and was attributed to the formation of complexes between the o-quinones and sulfite before the onset of browning (Sayavedra-Soto and Montgomery 1986). The inhibitor is gradually consumed in the quinone-coupling process: thus its action depends on its concentration, as well as on the nature and concentration of the phenols present (Vamos-Vigyazo 1981). In the presence of o-dihydroxy phenols, sulfite may get entirely consumed before complete enzymatic inactivation has occurred (Vamos-Vigyazo 1981). In such a case color formation is delayed and attenuated, but not entirely eliminated (Haisman 1974). In the presence of both mono- and dihydroxy phenols, ascorbic acids interferes (by its reducing action) with the quinone coupling to sulphite, and thus promotes sulphite inhibition of the enzyme (Vamos-Vigyazo 1981).

Sodium metabisulfite was the most potent inhibitor of Emir grape PPO with 5.5, 13.5, 41.1 and 100% inhibition with concentrations of 0.05, 0.1, 0.25 and 0.50 mmol/L respectively (Unal and Sener 2006). It was reported that sodium bisulfite has a direct irreversible inhibition effect on mushroom PPO with an I<sub>50</sub> value (concentration required to reduce the observed activity by 50%) of 0.20mM (Golan-

Goldhirsh and Whitaker 1984). Presence of sulfites in standardized foods should be declared on the label when the sulfating agents have a functional effect or are present at a detectable level, defined as 10 ppm or more (Title 21, U.S. Code of Federal Regulations 101.100). Use of sulfites to inhibit browning presents some disadvantages: 1) the corrosion of machinery 2) the destruction of nutrients, 3) the production of tissue softening and off-flavors, and 4) adverse health effects (Girelli and others 2004).

Sorbates are considered GRAS. The presence of sorbic acid as an antimicrobial agent had a minimal effect on the browning of avocado puree (Soliva-Fortuny and others 2002). Sorbates are unstable in aqueous solution and suffer oxidative degradation that depends on pH and the presence of other additives (Campos and Gerschenson 1996; Campos and others 1997). Autoxidative degradation of sorbates proceeds at higher rates at acidic pH values due to the fact that only undissociated molecules are prone to oxidative degradation (Arya 1980). Carbonyl compounds are formed during the degradation of sorbate that can take part in non-enzymatic browning reactions and lead to undesirable changes (Campos and others 1997).

Ascorbic acid is a highly effective inhibitor of enzymatic browning primarily because of its ability to reduce quinones to phenolic compounds before they can undergo further reactions to form pigments (Sapers 1993). However, once added, ascorbic acid has been completely oxidized to DHAA by this reaction, and quinones can accumulate and undergo browning and DHAA can brown non-enzymatically (Sapers 1993).


Since PPO is a metalloprotein with copper as the prosthetic group, it can be inhibited by metal chelating agents. Distinct differences were found in the action of this group of inhibitors upon PPO from chloroplasts, mitochondria or the soluble fraction (Vamos-Vigyazo 1981). PPO can also be inhibited by procyanidins, native and oxidized, and by the oxidation products of caffeoylquinic acid and (-)-epicatechin (Le-Bourvellec and others 2004). Some compounds like cyclodextrins have dual effects, activating or inhibiting browning, according to the plant material used (Sojo and others 1999). Proteins, peptides, and amino acids can effect PPO activities by reacting with o-quinones and by chelating the copper at the active site of PPO (Girelli and others 2004). Dipeptides vary widely in their inhibition effect on mushroom PPO ranging from direct, indirect or no inhibition at all (Girelli and others 2004). In recent studies, sodium chlorite emerged as a potential PPO inhibitor with mixed inhibition that is dependent on pH and inhibitor concentration (Lu and others 2006). The search for naturally occurring inhibitors has led to the discovery of a number of active compounds like chalcones and related compounds (Mayer 2006).

Oxalate in spinach leaves and quercetin and leucoanthocyanins in tea leaves have been described as natural inhibitors of catechol oxidase (Mayer 1987). PPO is inhibited by trans-rasveratrol with a linear relationship between PPO activity and trans-rasveratrol concentration (Fan and Mattheis 2001). Inhibition of apple PPO by native and oxidized procyanidins which are oxidation products of caffeoylquinic acid and (-) epicatechin, was reported by Le-Bourvellec and others (2004).

## **Anthocyanins**

Anthocyanins are the most important water-soluble pigments in plant tissue, and impart blue, red and purple colors. The anthocyanins (Greek *anthos*, flower and *kyanos*, blue) are part of the very large and widespread group of plant constituents known collectively as flavonoids. The basic structure of anthocyanins is a flavylum cation with different groups attached to different positions in the molecule. The six most common anthocyanidins or aglucones out of the 17 that occur in nature are: pelargonidin, cyaniding, peonidin, delphinidin, petunidin and malvidin (Figure 3). Differences between the individual anthocyanins are the number of hydroxyl groups in the molecule, the degree of methylation of these hydroxyl groups, the nature and number of sugars attached to the molecule and the position of attachment, and the nature and number of aliphatic or aromatic acids attached to the sugar in the molecule (Mazza and Brouillard 1987b). The sugars most commonly attached to anthocyanidins are glucose, galactose, rhamnose and arabinose (Mazza and Brouillard 1987b). Sugar present in anthocyanin molecule imparts a higher solubility and stability. Sugars, acylated sugars, methoxyl and hydroxyl groups have a marked effect upon the color and reactivity of anthocyanins (Mazza and Brouillard 1987b). Fifteen anthocyanins have been identified in the blueberries cultivated in North America and they are 3-arabinoside, 3-galctoside, and 3-glucoside of peonidin, cyaniding, malvidin, delphinidin and petunidin (Sapers and others 1984; Ballington and others 1987; Mazza and Miniati 1993). The color of anthocyanin-containing media depends on structure and concentration of the pigment, pH, temperature, presence of copigments, metallic ions, enzymes, oxygen, ascorbic acid, sugars and

their degradation products, sulfur dioxide, actual anthocyanin concentration and other factors (Mazza and Miniati 1993). Generally as the number of phenolic hydrolysis increases, the color changes from pink to blue. Methoxyl groups replacing hydroxyl reverse the trend (Mazza and Brouillard 1987b). In aqueous media, most of the natural anthocyanins behave like pH indicators. Anthocyanin solutions are red or yellow at low pH's in a form of a flavylum cation, blush at intermediate pH's in the form of a quinoidal base, and colorless at high pH in a form of carbinol pseudobase and chalcone form (Mazza and Brouillard 1987a).



Name	R <sub>1</sub>	R <sub>2</sub>
Delphinidin	OH	OH
Petunidin	OCH <sub>3</sub>	OH
Cyanidin	OH	H
Pelargonidin	H	H
Peonidin	OCH <sub>3</sub>	H
Malvidin	OCH <sub>3</sub>	OCH <sub>3</sub>

Figure 3. Structure of most commonly found anthocyanidins.  
(<http://health.wedar.com/show.asp?id=4701>)

It is generally accepted that in acidic solutions anthocyanins exist as equilibrium mixtures of the colored flavylum cation and the colorless carbinol or pseudo base (hemiketal form) (Timberlake and Bridle 1967b) (Figure 4). Thus for example for cyanidin 3-glucoside, at a pH of 3.01, 50 % will be in the colored form and 50% will be colorless (Wrolstad 2004).




Figure 4. Forms of anthocyanins present in acidic solution

The electron deficient anthocyanins molecule is susceptible to nucleophilic attack of compound which may exist naturally in the plant and food materials. Foremost among these nucleophiles is ascorbic acid, which has been shown to accelerate anthocyanin breakdown. Anthocyanins can disappear as monomeric compounds and transform into polymeric forms which result in a color change to a brown pigmentation (Iversen 1999).

The anthocyanins pigment of the native European blueberry or bilberry (*Vaccinium myrtillus*) have long been used for improving visual acuity and treating

circulatory disorders (Mazza and Miniati 1993). There is experimental evidence that certain anthocyanins and flavonoids have anti-inflammatory properties, and there are reports that orally administered anthocyanins are beneficial for treating diabetes, ulcers and may have antiviral and antimicrobial activities and can also provide protection against UV radiation (Mazza and Miniati 1993). Anthocyanins, as a functional food component, can aid in the prevention of obesity and diabetes (Tsuda and others 2003). The chemical structure (position, number and types of substitution) of an individual anthocyanin molecule has a bearing on the degree to which anthocyanins exert their bioactive properties (Lila 2004). The nature of the sugar conjugated and the aglucone are important determinants of anthocyanin absorption and excretion in both human and rats (McGhie and others 2003). Delphinidin but not malvidin or cyanidin, provided endothelium-dependent vasorelaxation in the rat aorta (Andriambeloson and others 1998).

### **Degradation of anthocyanins by PPO**

It is well known that anthocyanins are poor substrates for PPO, and this is probably due to the presence of the sugar moiety that causes steric hindrance, since the aglucon forms are often oxidized by PPO (Mathew and Parpia 1971). It has been reported that PPO can act on anthocyanins when triphenolic function is present on the B ring of the flavylum structure (delphinidin derivatives) (Sakamura and Obata 1963; Sakamura and others 1965). The loss of about 20% of pigments in a model containing PPO and cyanidin seems to indicate that the enzyme could act directly on this pigment (Wesche-Ebeling and Montgomery 1990b). This could be due to the

presence of cyanidin in the aglucone (no sugar attached) form although the concentration of cyanidin in the aglucone form was not determined (Wesche-Ebeling and Montgomery 1990b). Enzymatic oxidation of phenolic compounds that are present in blueberries by PPO can occur during processing as soon the berries are damaged. The first reaction step that occurs is the oxidation of phenolic substrates to their o-quinones. Quinones are very reactive species which are able, as oxidants, to oxidize other substrates, being reduced in the process to the original phenol, and as electrophiles to react with various nucleophiles (Sarni and others 1995). These o-quinones reactions, namely oxidation and condensation reactions can lead to discoloration and can involve other phenolic compounds, especially anthocyanins. The use of model systems has shown that the enzymatically generated o-quinones play an essential role in the process of anthocyanin degradation (Kader and others 1999a). Although anthocyanins are not direct substrates for PPO, anthocyanins with an o-diphenolic B ring were enzymatically oxidized by generated o-quinines of catechol, catechin, chlorogenic acid (Peng and Markakis 1963; Pifferi and Cultrera 1974; Raynal and Moutounet 1989; Wesche-Ebeling and Montgomery 1990b) and caftaric acid (Cheynier and others 1994). Degradation of anthocyanins occurs when active PPO and its substrate are present in the system. In the model system containing chlorogenic acid, the degradation rate of PPO and anthocyanins was much faster than without chlorogenic acid, and no degradation occurred when the enzymatic extract was inactivated by heating and chlorogenic acid was also not oxidized (Kader and others 1997a). Degradation of anthocyanins is also tied to its structure, which is strictly correlated to the reaction pH (Pifferi and Cultrera 1974).

Whatever the PPO substrate used, the degradation of anthocyanins increased as the pH increased with the sharp rise at pH 4.2 (Pifferi and Cultrera 1974). Sarni-Manchado and others (1997) showed that a hemiacetal form of malvidin are more reactive than the flavylium form which is directly dependent on pH. Mechanism and rate of anthocyanin degradation by o-quinones is closely related to their structure, especially their B ring. Sarni and others (1995) concluded that the first reaction step that leads to the discoloration of grape must like model solution depends on the nature of the anthocyanin: o-diphenolic anthocyanins behave mostly as reductants whereas non o-diphenolic anthocyanins act as nucleophiles to yield condensation products. These researchers also concluded that in complex media like grape must, both reactions occur simultaneously but o-diphenolic anthocyanins are degraded faster.

In the presence of chlorogenic acid and enzyme extract, cyanidin-3- rutinoside was rapidly degraded and could not be detected after 20 min of reaction (Raynal and Moutounet 1989). The quinones that are formed from the action of PPO on chlorogenic acid seem to be responsible for the oxidation of anthocyanins because degradation was not evident when quinones were blocked by glutathione (Raynal and Moutounet 1989). In the oxidizing grape must concentration of o-diphenolic anthocyanins like delphinidin cyanidin and petunidin decreased much faster than non o-diphenolic anthocyanins like malvidin and peonidin (Cheynier and others 1994). Degradation of cyanidin- glucoside (o-diphenol) was much faster than the degradation of pelargonidin-3-glucoside and proceeded by a mechanism of coupled oxidation in the presence of the chlorogenic acid o-quinone (Kader and others 2002). The secondary anthocyanin o-quinones generated by coupled oxidation also

proceeded further to condensation products (Sarni and others 1995). Once formed, all condensation products were rapidly degraded to a colorless compound, either by enzymatic oxidation or by reaction with primary or secondary quinines (Sarni and others 1995).

The mechanism of anthocyanin degradation depends not only on the structure of the anthocyanin but also whether the degradation was monitored on one particular anthocyanin or a mixture of anthocyanins. Cheynier and others (1994) reported that non-o-diphenolic anthocyanins may react with caffeoyltartaric o-quinone or secondary quinones that are formed from delphinidin and petunidin-3-glucoside leading to the formation of copolymers. These reactions have also been mentioned by Sarni and others (1995) and can only take place in a mixture of o-diphenolic and non-o-diphenolic anthocyanins.

### **Degradation of o-diphenolic anthocyanins**

Cyanidin glycoside is a diphenolic anthocyanin that is very susceptible to a reaction with o-quinines, secondary products of phenolics oxidation by PPO. Sarni and others (1995) proposed a mechanism of cyaniding-3-glucoside by PPO in the presence of caffeoyltartaric acid o-quinine generated by PPO in the presence of caffeoyltartaric acid as a substrate in a 2 stage process:

1. caffeoyltartaric acid o-quinine + o-diphenolics anthocyanin = caffeoyltartaric acid + anthocyanin o-quinine

Quinines of cyaniding-3-glucoside are extremely unstable and proceed readily to condensation products :

2. anthocyanin o-quinone + anthocyanin = caffeoyltartaric acid-anthocyanin product

The addition of caffeic acid quinone to a cyanidin-3-glucoside solution resulted in the disappearance of the red color and the concomitant formation of a slight orange color (Kader and others 1999a). Kader and others (1999a) proposed identical mechanism of degradation for cyanidin-3-glucoside with caffeic acid quinone as Sarni and others (1995). They did not observed any brown color in the reaction mixture, which means that degradation products of cyanidin-3-glucoside are not involved in the formation of brown polymers. Kader and others (1998) incubated equimolar concentrations of chlorogenic acid and cyanidin-3-glucoside with blueberry PPO and reported that after 10 min of reaction, pigment could not be detected in the reaction mixture and after 20 min, the reaction turned brown. These researchers concluded that cyanidin-3-glucoside is oxidized by a coupled oxidation mechanism involving chlorogenic acid o-quinone that is generated by PPO in the presence of chlorogenic acid which leads to complete discoloration. The ratio of degraded cyanidin-3-glucoside to oxidized chlorogenic acid is relatively constant and equal 2, which means that part of the chlorogenic acid is incorporated into degradation products of cyanidin-3-glucoside (Kader and others 1998)

### **Degradation of non-diphenolic anthocyanins**

The degradation of non-o-diphenolic anthocyanins has been regarded as a two step process involving the enzymatic oxidation of the o-diphenolic substrate into the corresponding o-quinone, followed by the reaction of the enzymatically generated o-

quinone with the anthocyanins (Kader and others 1999b). The latter reaction leads to the formation of adducts which can be oxidized either by enzymatic oxidation or by reaction with the o-quinone with no coupled oxidation occurring as expected by their structure (Sarni and others 1995).

Malvidin glycosides are non o-diphenolic anthocyanins that are less susceptible to the degradation of PPO generated o-quinines. However, Sarni and others (1995) noticed a decrease in malvidin-3-glucoside in the model system in the presence of PPO and caffeoyltartaric acid (PPO substrate) and attributed that decrease to following reaction;

caffeoyltartaric acid o-quinine + anthocyanin = caffeoyltartaric acid –anthocyanin  
product

There was no degradation of malvidin-3-O-glucoside in the presence of PPO without the substrate (caftaric acid), and no new products appeared in the system with malvidin-3-O-glucoside and caftaric acid as a PPO substrate (Sarni-Manchado and others 1997). This indicated that malvidin-3-O-glucoside is not a direct substrate for PPO and that it does not directly interact with the PPO substrate, caftaric acid in this case (Sarni-Manchado and others 1997). They also concluded that at a pH of 3.4, (where 2 forms of malvidin-3-O-glucoside, flavylium and hemiacetal coexist) both reacted with caftaric acid o-quinone that was formed by the action of PPO forming colored flavylium/caftaric acid and hemiacetal/caftaric adducts.

Weische-Ebeling and Montgomery (1990) studied the degradation of pelargonidin-3-glycoside by strawberry PPO. A model system containing pelargonidin-3-glycoside + PPO showed a slight loss of pigment (5%), but in the

presence of catechin as a PPO substrate, 50% of pelargonidin-3-glycoside was lost after 24 h. The authors proposed a mechanism of degradation by incorporation of anthocyanins into condensation products of catechin by quinone-phenol reaction. Kader and others (1999) reported that pelargonidin-3-glycoside is not a substrate for PPO since no molecular oxygen was consumed in the mixture of PPO and pelargonidin-3-glycoside. The degradation of pelargonidin-3-glycoside occurred only when both chlorogenic acid and blueberry PPO were present in the mixture. These researchers suggested that chlorogeno-quinone that was generated during the oxidation of chlorogenic acid by PPO plays an important role in the mechanism of pelargonidin-3-glycoside degradation and that the simplest hypothesis to explain this behavior is the concept of a condensation reaction between the chlorogeno-quinone or its degradation products and pelargonidin-3-glycoside. Pelargonidin 3-glucoside reacted with the hydroxycaffeic acid o-quinone formed by the dismutation of caffeic acid in aqueous media which led to the formation of condensation products that contained both caffeic acid and pelargonidin 3-glucoside moieties (Kader and others 2001)

### **Inhibition of anthocyanins degradation**

Degradation of anthocyanins can be inhibited by the direct inhibition of PPO and with that, formation of o-quinines or by the inhibition of reaction of anthocyanins with already generated o-quinones. The quinones formed from the action of PPO on chlorogenic acid seem to be responsible for the oxidation of anthocyanins because the degradation was not evident when quinones were blocked by glutathione (Raynal and

Moutounet 1989). Cheynier and others (1994) also reported that the degradation of anthocyanins was totally inhibited by the addition of glutathione in grape must prior to oxidation. In the conditions when glutathione was added, caffeoyltartaric acid quinones that were by enzymatic oxidation were converted to other products, which prevented them from reacting and degrading the anthocyanins. With larger amounts of the inhibitor glutathione, competition between trapping of the quinones by glutathione and their reduction by anthocyanins was in favor of the former (Cheynier and others 1994)

Addition of ascorbic acid as a reducing agent to the reaction mixture containing pelargonidin-3-glucoside, chlorogenic acid and blueberry PPO induced instantaneous bleaching after 2 min of reaction, (Kader and others 2001). These researchers concluded that in the presence of excess reducing agent, the o-quinones that were formed were reduced to original phenol. On the other hand Sakamura and others (1965) reported that the addition of ascorbic acid retards the loss of pigment as long as ascorbic acid is present in the mixture. Inhibition of PPO activity by phenylthiourea prevented oxidation of chlorogenic acid which in addition prevented the degradation of cyanidin-3-rutinoside (Raynal and Moutounet 1989).

Addition of sulfites was reported both to degrade anthocyanins and to inhibit their degradation by the action of PPO. This bleaching can be reversible or irreversible. The reaction of sulphites with anthocyanins, nicotinamide derivatives, the flavin moiety of flavoenzymes and folate all cause reversible binding of the additive (Wedzicha 1992). The reversible reaction with sulfur dioxide involves reaction with colored flavylium to form colorless chromen-2 (or 4)-sulphonic acid,

which is similar in structure and properties to an anthocyanin carbinol (pseudo) base (Jurd 1963). When sulphur dioxide was added to a diluted blackcurrant juice, the visible color fell rapidly to a minimum value and then rose slowly during the course of several hours (Timberlake and Bridle 1967a). This recovering of the color was attributed to the other substances present in juice (e.g. sugars and dehydroascorbic acid) that were competing with the anthocyanins for  $\text{SO}_2$  and were slowly removing it from the initially formed anthocyanin complex (Timberlake and Bridle 1967a).

Sulfur dioxide (8ppm) completely inhibited the degradation of anthocyanin in a model system containing mushroom PPO and catechol as a substrate at pH 6.5 (Goodman and Markakis 1965). In tart cherry juice under similar conditions, but with no catechol added, 30ppm of  $\text{SO}_2$  was required for complete inhibition of anthocyanin degradation (Goodman and Markakis 1965). These greater concentration of  $\text{SO}_2$  in the juice than in the model system that was necessary for the inhibition of PPO was attributed to the  $\text{SO}_2$ -binding capacity of carbonyl compounds that are present in the juice.

Influence of the added  $\text{SO}_2$  on the content of the colored anthocyanins depends of the content of the  $\text{SO}_2$ -binding carbonyl compounds present in wine with the same level of total  $\text{SO}_2$  that differently influences anthocyanin color in different wines (Dallas and Laureano 1994). Malvidin 3-glucoside is immediately decolorized by an excess of  $\text{SO}_2$  (2000mg/l) (Dallas and Laureano 1994). It was found that  $\text{SO}_2$  had a strong stabilizing effect on pelargonidin 3-glucoside in canned and bottled strawberries (Adams and Ongley 1973) and in strawberry juice (Bakker and Bridle 1992). Sodium sulphite was found to have a strong stabilizing effect on

pelargonidin 3-glucoside in canned and bottled strawberries, and it was originally theorized that this was either due to its antioxidant action or to the fact that it forms addition compounds with aldehydes such as furfural and 5-hydroxymethylfurfural (arising from the break down of sugars) which might otherwise condense with the anthocyanin to yield colorless or brown compounds (Adams and Ongley 1973). Incorporation of sulfur dioxide in the extraction medium significantly increased the yield of anthocyanins from apple peel (Timberlake and Bridle 1971).

### **Blueberry juice**

Most of the blueberries in Mississippi are sold on the fresh market ([http://www.mdac.state.ms.us/n\\_library/pub\\_form/publications/pdf/com\\_blueberries.pdf](http://www.mdac.state.ms.us/n_library/pub_form/publications/pdf/com_blueberries.pdf).) Mississippi's early season helps growers get top prices before northern blueberries start competing in the market. The perishable nature of blueberries in the fresh form coupled with the decrease in price in southern berries once northern production starts, requires rapid movement to market (Sadfar and Albert 2000) or further processing. There is also a lot of excess fruit that includes unripe, blemished and broken berries that are not suitable for the fresh or frozen markets. Frozen berries are difficult to sell profitably due to supply exceeding demand and high storage costs (Main and others 2001). Use of excess fruit in juice products is especially attractive since machine-harvested fruit can be used (Bakker and others 1998). Health attributes ascribed to blueberries create a need for the development of processed blueberry products that are convenient for the consumer. Of all fruits and vegetables, blueberries are ranked highest in their antioxidant activity (Prior and others 1998).

Both cranberry and blueberry juice contain high molecular weight constituents which selectively inhibit mannose resistant adhesions that are produced by urinary isolates of *Eschericia coli* by binding to the bacterial surface, possibly to the adhesion itself (Ofek and others 1996).

Blueberries are often pressed into juice and juice concentrate, to reach more wide spread consumption (Rossi and others 2003). Recent reports in the beverage industry indicate an increase in the production of functional beverages. Blueberry juice would be a rich source of antioxidants and could be classified as a functional beverage. Blueberries may be the biggest berry beverage breakthrough yet. In Asia and Europe, blueberries are one of the most popular flavors for fruit beverages and juices (<http://www.foodprocessing.com/articles/2005/559.html?page=2>). In North America, blueberries in beverages are suddenly becoming popular with companies like Izze Beverage Co., extending their line of pure fruit juice sodas with blueberry last year; Leading Brands released TrueBlue blueberry juice cocktail; Island Juice Company produces a line of berry blend juice beverages; Ocean Spray introduced its new cocktail, Organic Cranberry-Blueberry, for a double-shot of health (<http://www.foodprocessing.com/articles/2005/559.html?page=2>).

Blueberry juice is an expensive product because blueberries are priced higher than other small fruits (Main and others 2001). Blueberry juice can be also mixed with other juices and still produce a blueberry like juice or juice cocktail (Main and others 2001). Berry juices have become widely used in blends with the more traditional juices, such as apple, cranberry, and orange juice (Roberts and others 2004).

The percentage of blueberry juice in the blend and the type of blending juice greatly affects the final product. Concord and Venus grapes blended with blueberry juice resulted in a flavor similar in intensity to pure blueberry juice while apple and cranberry blends produced a juice with diminished blueberry flavor (Main and others 2001).

### **Blueberry juice processing methods**

Juice processing methods affect the quality of the final product. The stability of color in blueberry is an important aspect of controlling quality during processing and storage (Yang and Yang 1987). Anthocyanins exist almost exclusively in the skin, whereas phenolics and antioxidant properties are mostly in the flesh (Lee and Wrolstad 2004). Since blueberries contain anthocyanins that impart the red color in the skin, breaking the skin and maceration of the blueberry mash are necessary steps in order to extract the color and obtain good quality juice. Freezing and heating of the blueberries had a more pronounced effect than pectolytic enzymes on the skin cells, and consequently on the liberation of color pigments. The loss of semipermeability and mechanical injury to the cell membranes appears to determine the total color released (Fuleki and Hope 1964). Crushing of the berries and heating have been used to increase color extraction from the berries and increase yield. Fuleki and Hope (1964) compared three treatments; hot pressing (62°C for 30 min), cold enzyme pressing and hot enzyme pressing on blueberry juice yield and composition. They reported that the hot enzyme treatment was superior to the other two, although the higher extraction of color was mainly attributed to the increased

temperature and mechanical injury to the skin rather than the action of pectolytic enzymes. Lee and others (2002) also concluded that heating of berries may have contributed towards a breakdown of the skins, which could have increased color extraction, but still with a substantial amount of anthocyanins in the press cake. Reextracting the press cake or crushing the blueberries into fine particles may aid in greater extraction of anthocyanins into the final product (Lee and others 2002). Blueberry puree that was held for 60 min at 60°C had higher anthocyanin and phenolic concentration than puree held at 25°C, with anthocyanins being more affected than phenolics (Kalt and others 2000). The increased extraction of anthocyanins was attributed to the increased permeability of membranes in the macerated berries at higher temperatures, and to decreased solubility of oxygen at higher temperatures that decreases oxidative degradation (Kalt and others 2000). Skrede and others (2000) obtained 83% yield of juice with the press cake residue accounting for 10% of the starting material. They also had 7% loss of material in the milling, enzyme maceration and pressing unit operations. Processing operations for the production of juice included thawing (5°C), milling (6-7mm), depectinization at 43°C for 2h, pressing (0.5bar), pasteurization (90°C, 1 min), and filtering to obtain single strength juice with 15 Brix and further concentration at 40°C to 73.5 Brix. Lee and others (2002) compared two pretreatments: addition of 100 ppm SO<sub>2</sub> and heat (95°C for 2 min) followed by crushing, pectinase treatment, pressing, clarification, pasteurization (90°C for 90s) and concentration on juice quality. Yield of the juice ranged from 74 to 89% (w/w), with anthocyanins not efficiently extracted, which had a negative influence on juice quality. Pretreatments with heat and SO<sub>2</sub> resulted in

higher recovery of red color pigments with the more intense color (higher chroma) in pasteurized juice compared to control, while the heat treatment samples were darker (smaller L values) than the control and SO<sub>2</sub> treated juice (Lee and others 2002). Rossi and others (2003) reported a juice yield that ranged from 79 to 81% (w/w) when they used a process that consisted of thawing, blanching, milling, depectinization for 1h at room temperature with enzymes, pressing and pasteurization. They concluded that the addition of steam blanching of the fruits should be considered a very important factor when evaluating processed blueberry products for their possible health benefits (Rossi and others 2003). The cold processing of blueberries was not as efficient at transferring antioxidants or phenols to the final juice as methods where heat was used (Carlson 2003). Pasteurization using a steam kettle was the most effective at retaining antioxidant activity as measured by total phenols and ORAC in the final products. One likely reason for this significantly higher level of antioxidant retention is due to exposure to higher heating temperatures resulting in increased permeability of water soluble substances, which then diffused into the liquid stream. The higher heat also degrades enzymes (PPO and glucosidase) that would be harmful to the anthocyanins and lower the oxygen concentration in the final products (Carlson 2003).

### **PPO in food processing**

Polyphenol oxidase (PPO) is a very important enzyme in the food industry. Processing or any wounding of fruits and vegetables may cause cell disruption that can lead to quinone formation due to the action of PPO with subsequent product

deterioration. The appearance of food and beverages may be affected but also the taste and nutritional value, often decreasing the quality of the final products (Gandia-Herrero and others 2005). For the fruit and vegetable processor, the action of PPO is primarily connected to the enzymatic browning of fresh and off-flavor generation in canned or frozen horticultural products, respectively (Vamos-Vigyazo 1981). Both phenomena are of vital importance to the manufacturer as they impart not only the sensory properties and hence, the marketability of a product, but often lower its nutritional value (Vamos-Vigyazo 1981). Enzymatic oxidation of phenolic compounds present in blueberries by PPO can occur during processing as soon the berries are damaged. The first reaction step that occurs is oxidation of phenolics and conversion of substrates to their o-quinones. Quinones are very reactive species which are able, as oxidants, to oxidize other substrates, being reduced in the process to the original phenol, and as electrophiles to react with various nucleophiles (Sarni and others 1995). These reactions of o-quinones, namely oxidation and condensation reactions can lead to discoloration and can involve other phenolic compounds, especially anthocyanins. Degradation of anthocyanins occurs when active PPO and its substrate are present in the system. Although anthocyanins are not direct substrates for PPO, anthocyanins with an o-diphenolic B ring were oxidized via enzymatically generated o-quinines of catechol, catechin, and chlorogenic acid (Peng and Markakis 1963; Pifferi and Cultrera 1974; Raynal and Moutounet 1989; Wesche-Ebeling and Montgomery 1990b) and caftaric acid (Cheynier and others 1994). Thermal treatment and addition of chemical inhibitors are some ways to inhibit the action of PPO.

PPO is not an extremely heat-stable enzyme. Short exposure, in the tissue and solutions, to temperatures of 70 to 90°C are, in most cases, sufficient for partial or total irreversible destruction of its catalytic function (Vamos-Vigyazo 1981). Blanching (3 min in a steam blanching tunnel) of blueberry fruit was extremely effective at reducing PPO activity, maximizing anthocyanin recovery in the juice (Rossi and others 2003). The PPO in crude extracts of plum, apple, pear and avocado was inactivated at 60-65°C, whereas for grape PPO, a temperature of 55°C was enough for inactivation (Weemaes and others 1998). PPO thermal stability can be affected by some substances that are present in the food (Tomas-Barberan and Espin 2001a). Addition of both EDTA and benzoic acid increased the thermal stability of mushroom PPO, whereas glutathione produced a sensitization to temperature treatments, probably due to an interaction with a disulphide bond of the enzyme (Weemaes and others 1997).

It has been reported that heat inactivation of PPO is unacceptable in anthocyanin-containing juice products, since the high temperature required for PPO inactivation can cause anthocyanin degradation (Siddiq and others 1992). Carboxylic acid is a strong inhibitor of PPO (Pifferi and others 1974; Walker and Wilson 1975; Walker and McCallion 1980; Gunata and others 1987; Ferrar and Walker 1996). Benzoic acid and some derivatives of cinnamic acid are competitive inhibitors of PPO (Pifferi and others 1974). When 4-methyl catechol is used as a substrate, benzoic acid is a competitive inhibitor of PPO (Gunata and others 1987; Janovitz-Klapp and others 1990b). Benzoic acid at a concentration of 2.5mM (0.03%), inhibited 43% of grape PPO at pH 5 (Gunata and others 1987). The presence of 5mM benzoic acid

lowered enzyme activity by 5%. At higher concentrations of benzoic acid, the enzyme activity was reduced more: 11% at 10mM, 36% at 25mM and 52% at 50mM (Weemaes and others 1997). This decrease in activity is probably due to competition between benzoic acid and catechol for the active site of the enzyme (Weemaes and others 1997).

### **Acids in blueberries**

Sixteen different organic acids were identified in highbush blueberries with the predominant acids being citric, malic, quinic and chlorogenic, averaging 70%, 7%, 4% and 16%, respectively (Markakis and others 1963). In rabbiteye fruit, the percentage contribution by citric, succinic, malic and quinic acid is about 10%, 50%, 34% and 6%, respectively (Kalt and others 1996). Other researchers reported 95% citric acid and 1% to 2% each of quinic and malic acid in ripe Wolcott fruit (Kushman and Ballinger 1968). Succinic and malic acids were the predominant acids in rabbiteye cultivars averaging 50% and 33%, respectively (Ehlenfeldt and others 1994). Among rabbiteye clones, citric acid averaged 10% and was never found to be more than 22%, while quinic acid was consistently present as only a minor constituent, averaging 6% (Ehlenfeldt and others 1994). Quinic acid is found in plants and microorganisms and has a regulation role in the biosynthesis of aromatic compounds in the shikimate pathway (Dewick 1998). The biosynthesis of quinic and shikimic acid are interlinked and both are targets in the search for the new herbicidal, antifungal, antibacterial and antiparasitic agents that may not affect mammals (Pansare and Adsool 2006). High concentrations of quinic acid, between 0.5 and 1.18%, were

reported in juices from different varieties of cranberries and lingonberries but only trace amounts were found in blueberries from Germany and Argentina (Jensen and others 2002). Quinic acid is also starting material in the synthesis of viral neuraminidase inhibitors for the treatment of influenza (Kim and others 1997). Quinic and shikimic acids are gaining interest because they can be a starting material in the production of the bird flu medicine, Tamiflu (Bradley 2005). Shikimic acid is also a starting material for the making of Tamiflu, but there is only a limited supply of the acids, with the most abundant source from star anise and the leaves of ginkgo biloba (Bradley 2005). Tamiflu can also be synthesized from quinic acid found in cinchona bark, but supplies are also limited (Bradley 2005).

## **Phenolics**

Phenolics are present in fruits, vegetables, leaves, nuts, seeds, flowers and barks. Although structurally diverse, phenolics can be classified into two groups – the flavonoids and the nonflavonoids. The flavonoid family includes flavonols (myricetin, quercetin, kaempferol and isorhamnetin), flavan-3-ols (catechin and epicatechin) and anthocyanins. The nonflavonoids encompass gallic acid, hydroxycinnamates (p-coumaric acid, caffeic acid, and ferulic) and resveratrol. Anthocyanin flavonoids and hydroxycinnamates are the largest constituents of the total phenolic content of blueberries (Kalt and others 2000). Blueberries are very rich in total phenolics. The total phenolics content of highbush and lowbush blueberries is about 4-fold higher than in strawberries and raspberries (Kalt and others 1999). Reported values of total phenolics in rabbiteye blueberries vary between authors.

Values of 717-961 mg of gallic acid equivalents in 100 g of rabbiteye blueberries (Moyer and others 2002), 230.8 to 457.5 (Prior and others 1998), and 270.02 to 929.62 (Sellappan and others 2002), were reported depending on cultivar. Large differences in total phenolics within the same variety can be attributed to maturity stage of berries, weather conditions during the year, growing season, plant disease, geographic location, extraction method and many other factors (Sellappan and others 2002). Total phenolics play an important role in the plant tissue and the human diet. Many distinctive development features of fleshy fruits, such as loss of astringency and color, are related to changes in the synthesis and accumulation of phenolic compounds. Phenolic compounds are considered nonnutrient biologically active compounds (Sellappan and others 2002). The functionality of these compounds is expressed through their action as an inhibitor or an activator for a large variety of mammalian enzyme systems, and as metal chelators and scavengers of free oxygen radicals (Sellappan and others 2002). Several research groups have suggested the significance of fruit and vegetables phenolics as dietary antioxidants. Their studies suggested that phenolics have substantial antioxidant capacity and may reduce the risks of cardiovascular disease and cancer (Kalt and others 1999).

### **Changes in anthocyanins during juice processing**

Attractive color is a very important sensory characteristic for the consumer of fruits and processed fruit products (Garzon and Wrolstad 2002). Huge economic losses can also be accrued by the company due to the loss of attractive product color. A concord grape juice producer reported an approximate loss of \$ 700,000 due to the

loss of juice color (Wrolstad and others 1994). Not just because they impart color, but also because of their possible beneficial effect, particular attention has to be paid to the changes that anthocyanin pigments undergo with processing. Pronounced changes in anthocyanins occurred during all processing steps, with the most dramatic difference between the berry fruit and the initial pressed juice (Skrede and others 2000). Anthocyanins are not as efficiently extracted in the pressing operation as sugars, acids and other water solubles, which can have a negative impact on juice quality (Skrede and others 2000; Lee and others 2002). There was greater than 76% loss of anthocyanins in pasteurized juices when compared to frozen fruit (Lee and others 2002). Anthocyanins as well as other polyphenolics are readily oxidized because of their antioxidant properties and, thus, susceptible to degradative reactions during various unit processing operations (Skrede and others 2000; Rossi and others 2003). The oxidation of anthocyanins may be most significant in fruit juices or beverages, because the aqueous matrix can dissolve large amounts of oxygen during processing, as compared to products with low water content (Kalt and others 2000). The primary steps of processing (thawing, crushing, depectinization, and pressing) contributed a large loss in total anthocyanins (Skrede and others 2000; Lee and others 2002). It was shown that substantial losses of anthocyanins and other polyphenolics occurred when blueberries were processed into juice and that different classes of compounds had varying susceptibility to degradation, with different processing operations with the highest losses occurring during milling and depectinization due to the action of native PPO (Skrede and others 2000).

Stability of anthocyanins in food system is influenced by several factors, including chemical structure of the pigment, total anthocyanin concentration, presence of other phenolics and ascorbic acid, and processing conditions. The relative anthocyanins stability is related to their chemical structure. Diglycosidic substitution is reported to give more stability to the molecule than monoglycosidic substitution ((Mazza and Miniati 1993; Garzon and Wrolstad 2002). The delphinidin glycosides with the greatest lability have 3 ortho phenolics groups in the B ring and the cyanidin and petunidin derivatives, which have the second order of reactivity, have 2 orto phenolics groups (Skrede and others 2000). Peonidin and malvidin glycosides, which have the least reactivity, possess 1 phenolic substituent in the B ring with 1 and 2 adjacent metoxy substituents, respectively (Skrede and others 2000). Anthocyanin stability in products can also be related to total anthocyanin concentration in the products. The higher the total pigment concentration, the higher the stability of the berry juice or products (Garzon and Wrolstad 2002; Rein and Heinonen 2004). Color stability is more dependent on the total anthocyanin content rather than the qualitative anthocyanin composition (Skrede and others 2000). Lower pH (3.0) was found to stabilize the color of anthocyanins in the blueberry puree, because the equilibrium between colored flavylum and colorless pseudobase is shifted toward the flavylum at this pH (Yang and Yang 1987). Phenolics, especially phenolic acids have a protective effect on anthocyanin color (Rein and Heinonen 2004). Total anthocyanins concentration and ascorbic acid are also believed to be major factors that influence the stability of anthocyanins (Garzon and Wrolstad 2002). Fortification of syrups and jams with ascorbic acid caused a decrease in

pigment stability (Garzon and Wrolstad 2002). Not all anthocyanins are affected the same way during processing. Malvidin glycosides comprised 44% of the total anthocyanins in blueberry fruit and increased to 63% in pasteurized juice and concentrate (Skrede and others 2000). Similar results were reported by Lee and others (2002), in which malvidin glycosides in the berry were 51% of total anthocyanins and increased to 60-77% in pasteurized juice and concentrate. Delphinidin glucoside were the most unstable decreasing from 12% in blueberry fruit to 5% in pasteurized juice with only a trace amount present in the initial pressed juice (Skrede and others 2000). Decrease of delphinidin glycosides was also reported by Lee and others (2002) but only for the juice that received no pretreatment prior to pressing.

During juice processing, anthocyanins can be lost due to enzymatic breakdown, heat treatment and some are removed with the pulp (Iversen 1999). Activity of PPO is one of the most important factors that influence the color loss of Concord grape juices (Cash and others 1976). After addition of crude PPO extract to juice samples, formation of the brown precipitate was noticed and became greater as reaction time increased. The precipitate appeared to be the end result of enzymatic hydrolysis of anthocyanins to brown degradation products which then became polymerized and settled out of solution (Cash and others 1976). Chlorogenic acid will stimulate PPO destruction of anthocyanins during grape juice processing (Yokotsuka and Singleton 1997). Endogenous enzymes in blueberry fruit can cause pigment degradation in juice processing (Skrede and others 2000). The anthocyanin content of pasteurized single-strength juice was significantly higher than that of initial

pressed juice which was attributed to the action of the endogenous polyphenol oxidase (Skrede and others 2000; Lee and others 2002). Iversen (1999) also reported a higher anthocyanin content in the pasteurized black currant juice than in the raw juice. Kader and others (1997) also reported that endogenous PPO from highbush blueberries caused anthocyanin degradation in crushed fresh berries. The chlorogenic acid present in blueberries is an additional compositional factor which can contribute to the pronounced color degradation in blueberry juice (Skrede and others 2000).

Use of commercially available enzymes is widely used in the juice industry. Although most of the commercial enzyme preparations increase the yield of the juice, they also cause changes in the anthocyanin profiles. These changes include a decrease in anthocyanidin glycosides and formation of unknown components that are attributed to the high galactosidase, glucosidase and arabinosidase activities found in the enzyme preparations (Buchert and others 2005). Wightman and Wrolstad (1995) reported that destruction of the pigment most likely involves two steps. In the first step an enzymatic hydrolysis of the anthocyanin to anthocyanidin and sugar occurs, and then spontaneous transformation of the aglucone pigment causes the formation of colorless juice. As a result of all enzymatic treatments, practically no anthocyanidin galactosides were detected (Buchert and others 2005). Treatment without enzyme addition gave the highest yield of anthocyanins and phenolics (Lee and Wrolstad 2004).

Blanching of strawberries before juice and concentrate production had a protective effect on anthocyanin pigments, leucoanthocyanins, flavanols, total phenolics and ascorbic acid and also resulted in improved color stability (Wrolstad

and others 1980). Heating of the grape must before pressing had the greatest impact on the juice yield, amount of press fraction, pH, TA and color components, and nutraceutical analysis showed higher anthocyanin and phenolic content in the heated must (Threlfall and others 2005). After 2 h mash treatment at 50°C, 97% of the original anthocyanins content in the black currant berries were intact (Iversen 1999). Blanching of blueberry fruit (steam for 3 min) before milling induced higher anthocyanin retention (23 % instead 12%) when processed into juice with total anthocyanin content of juice from blanched blueberry twice the non blanched one (Rossi and others 2003). They concluded that better retention of anthocyanins for the blanched fruit could be due to the total inactivation of native PPO and increased skin permeability of fruit caused by the heat treatment. Rossi and others (2003) also observed that juice from blanched blueberry fruit before milling was more blue and less red than the juice obtained from the traditional process (without blanching). These color findings were in accordance with the anthocyanin recovery. Percent recovery in the blanching process of delphinidin glycosides and petunidin glycosides (tri-substituted), which are the most intense blue pigments, was higher than that of cyaniding-glycosides and peonidin glycosides (di-substituted).

Chemical treatments like SO<sub>2</sub>, citric acid and SnCl<sub>2</sub> had a stabilizing effect on strawberry anthocyanins during the processing of strawberry jam (Sistrunk and others 1982). Heat and SO<sub>2</sub> pretreatment before pressing the juice increased recovery of anthocyanins in all processing steps (Lee and others 2002). Less than 22% of the frozen berries' anthocyanins were present in the pressed juice with the control pressed juice having the lowest recovery when compared to the heat and SO<sub>2</sub> treated fruit

(Lee and others 2002). Heated and SO<sub>2</sub> treated pasteurized juices had 1.8 times the anthocyanin content of the control pasteurized juice (Lee and others 2002). Control pasteurized juice had a higher amount of anthocyanins than the initial pressed juice which was mainly attributed to enzymatic losses (Lee and others 2002). The press cake residues contained a substantial amount of anthocyanins (about 43-55% of frozen berry anthocyanins) with the control press cake containing the greatest amounts (Lee and others 2002). Heating of the berries may have contributed toward a breakdown of the skin, which would have increased the extractability of the anthocyanins (Lee and others 2002). Rossi and others (2003) demonstrated the effect of the blanching step prior to the milling of blueberries on the recovery of different anthocyanins is anthocyanin dependent. The anthocyanins that showed the highest benefit from the blanching treatment were, in decreasing order, the glycosides of delphinidin, petunidin, and cyanidin with the exception of cyanidin 3-glucoside that showed the lowest recovery (Rossi and others 2003). Malvidin and petunidin glucoside, having a single hydroxyl group on the phenolics ring were least affected by blanching and PPO inactivation (Rossi and others 2003). The highest percent recovery increase observed for delphinidin-glucoside could also be linked to their higher water solubility, due to the presence of three free phenolics functions (Rossi and others 2003). The proportion of malvidin-glycosides increased with the initial pressing compared to the fresh fruit. Malvidin-glycosides in the berry fruit were 51% of the total anthocyanins and increased to 60 to 77% in pasteurized juice and concentrate (Lee and others 2002). There was a concomitant decrease in delphinidin and petunidin glycosides (Lee and others 2002). Proportion of delphinidin-glycosides

decreased, especially in the control sample with only 8% left, while heat and SO<sub>2</sub> treated had 20 and 23% retained in the pasteurized juice (Lee and others 2002). The relative ratio of delphinidin glycosides versus cyanidin glycosides remained about the same in the juices as compared with the whole berry (Buchert and others 2005).

### **Changes in phenolics during juice processing**

Processing and storage can have a marked effect on the phenolic content of fruits, and might in turn affect their health promoting properties (Zafrilla and others 2001). As a general rule, during processing, phenolic biosynthesis is interrupted by the enzyme's destruction and/or cell structure degradation (Tomas-Barberan and Espin 2001a). Processing can enhance phenolic compound degradation (chemically or enzymatically if the oxidative enzymes have not been inactivated) or can produce chemical changes that affect quality characteristics (Tomas-Barberan and Espin 2001a). Lee and others (2002) reported considerable loss of phenolics during thawing, crushing and pressing with similar results obtained by Skrede and others (2000). Sixty to 65% of blueberry polyphenolics was lost during thawing, crushing and pressing (Lee and others 2002). There were no significant differences in polyphenolics taken after each processing step between the control, heat and SO<sub>2</sub> treated berries (Lee and others 2002). The press cake held 15 to 25% of the frozen berry polyphenolics, which is less than the proportion of anthocyanins left in the press cake (Lee and others 2002). Thirty six to 39% of the polyphenolics in the berries was present in the pasteurized juice. Forty two to 45% of the frozen berry phenolics was lost during juice processing and not accounted in the final pasteurized juices and

press cake (Lee and others 2002). Control pressed juice (no pretreatment) and SO<sub>2</sub> treated pressed juice had a lower value of polyphenolics than their pasteurized juices, but that was not the case for heat treatment. Hot pressed blueberries had higher amounts of total phenolics than cold pressed ones (Lee and others 2002).

Not all phenolic groups are affected the same by processing. Rossi and others (2003) observed a significant recovery of cinnamates when blueberry juice was blanched before milling, but lower than total anthocyanins. A significant increase in total cinnamates (phenolic) was observed in the heat treated juice, but lower than that of total anthocyanins (Rossi and others 2003). Extraction of cinnamates from the pulp is less affected by the heat treatment than that of anthocyanins from the fruit skin (Kalt and others 2000). Lee and others (2002) observed no changes in cinnamic acids of juices due to the heat treatment. Skrede and others (2000) reported a considerable loss of flavonol glycosides with processing with the pattern somewhat different than that of anthocyanins and chlorogenic acid. Some flavonol glycosides remained in the press cake residue (7%), but unlike with anthocyanins and chlorogenic acid, initial pressed juice contained higher levels than pasteurized single strength and concentrate which implied that flavonol glycosides were not as susceptible to enzymatic degradation as the anthocyanins and chlorogenic acid. Lee and others (2002) reported that higher levels of flavonol glycosides remained in the press cake (26-35% of the starting material's flavonol-glycosides), with just minor changes after the pressing step. An extensive decrease in procyanidin levels occurred during processing. The levels of procyanidin in initial pressed juice and single strength pasteurized juice was about 40% of the level in the fruit (Skrede and others 2000).

Thermal processing of raspberries into jam did not have much effect on the major phenolic present (Zafrilla and others 2001).

### **Non-thermal processing**

There is the demand of consumers and retailers for minimum processed foods but being safe at the same time. Therefore, producers and researchers are searching for the new food processing methods to gently preserve foods (Houska and others 2006). Even though processed blueberry products may retain a significant portion of antioxidant activity and total phenolic content as compared to fresh and frozen fruit, other forms of bioactivity like antiproliferation power may be compromised. Heat treated products suffered significant losses in antiproliferative activity, even in products in which total phenols and in vitro antioxidant activity were maintained (Schmidt and others 2005). Although thermal preservation methods provide safer foods, there is a loss in food quality that is associated with this processing method. Hence, the main objective of non-thermal preservation methods is to minimize the degradation of food quality through limiting heat damage of food (Senorans and others 2003). Compared with thermal processing, nonthermal methods provide high retention of flavors and nutrients, giving products a more natural taste (Zhong and others 2005).

### **Pulsed electric field**

Consumer demand for a higher quality, fresh appearing and safe food supply is the ultimate catalyst for the emergence of PEF on a commercial scale (Yeom and

others 2002). Thermal pasteurization is quite efficient in preventing microbial spoilage, but the applied heat may also cause undesirable nutritional and biochemical changes that can affect the overall quality of the final products. High voltage pulsed electric field (PEF) treatment is a promising non thermal processing technique that may radically change liquid food preservation technology. Treating liquid foods with PEF may inactivate microorganisms and enzymes with only a small increase in temperature, simultaneously providing consumers with safe, nutritious, and fresh like quality foods (Aguilar-Rosas and others 2007). PEF treatment might be suitable for the reduction of microbial cell counts in media that contain thermosensitive components like proteins and vitamins, which are difficult to pasteurize by heat processing (Grahl and Markl 1996). Reduction of living cell counts by PEF treatment in media that contain primarily yeast cells, i.e. fruit juices or other acid media, is of great interest, because in this case sufficient lethal effects are already produced by a very low energy output (Grahl and Markl 1996). Application of a pulsed electric field is restricted to food products that can withstand high electric fields, have low electrical conductivity, and do not contain or form bubbles (Senorans and others 2003). The products are minimally affected by the process since damage occurs on a cellular level and flavor and enzyme activity are not significantly diminished (Senorans and others 2003). Products with low pH's (such as apple juice, cranberry juice and orange juice) are more suitable for preservation by the PEF process than the neutral pH products such as milk. It is possible that the high acid environment of the food products may prevent sub-lethally damaged cells from recovery (Wouters and others 2001). The fact that PEF processes that are currently employed can not

inactivate bacterial spores limits the applications of PEF. As a consequence, only high acid conditions that can prevent outgrowth with a limited shelf life are applicable to preserve food products with this technology at present (Wouters and others 2001). PEF can possibly provide a reduction in the amount of energy that is required for fruit juice processing (Toepfl and others 2006).

The results of many researchers indicate that PEF operating with conditions with electric fields between 15 and 80kV/cm, pulse widths between 1 and 100Ms and the frequency of 1-100 pulses were sufficient for inactivation of bacteria and some certain enzymes, while taste and flavors are unaffected (Barbosa-Canovas and others 2001; Wesierska and Trziszka 2007). The initial counts of viable cells in cranberry juice were only around 100 cfu/ml and were not detected after PEF treatment (Jin and Zhang 1999). No differences were observed in the reduction of cell counts between thermal treatment and PEF treatment with an electric field strength of 40 kV/cm for a treatment time of 150 $\mu$ s (Jin and Zhang 1999). PEF treatment of cranberry juice using a bench scale PEF system at 20 kV/cm for 150 $\mu$ s resulted in about a 5 log cycle reduction in aerobic bacteria, yeast and molds, which was as effective as thermal treatment at 90°C for 90 min (Jin and Zhang 1999).

High intensity pulsed electric field (PEF) processing involves the application of pulses of high voltage (typically 20-80 kV/cm) to food that is placed between 2 electrodes. The PEF system usually consists of a high voltage pulse generator, a treatment chamber and a fluid handling system (Yeom and others 2002). The pulsed generator is capable of converting low voltage electricity into high voltage energy to be stored into capacitors until discharged. Electric pulses are generated when a pair

of high voltage electrodes are charged and discharged in a fraction of a second (Wesierska and Trziszka 2007). To process a product using PEF in a continuous system, the product flows through a series of PEF treatment chambers where it is exposed to the desired electric field strength for a desired amount of time (Yeom and others 2002).

It was observed that PEF-pretreatment caused an increase in anthocyanin concentration in resulting grape juice (Knorr 2003). PEF treatment caused the degradation of cyanidin-3 glycoside in aqueous-methanol solution. As the electric field intensity and the treatment time of PEF increased, degradation of cyanidin-3 glycoside increased significantly (Zhang and others 2007).

Organic acid and volatile aroma compounds of citrus juices did not change with PEF treatment of 28 kV/cm with 50 pulses (Cserhalmi and others 2006). When the most extreme PEF treatment was applied to orange juice, the decrease in the concentration of carotenoids with vitamin A activity was very small (Cortes and others 2006). PEF-treated orange juice retained better color and a higher concentration of vitamin C than heat-pasteurized orange juice through storage but no differences were found in pH, acidity and Brix (Elez-Martínez and others 2006). Antioxidant activities determined with DPPH method were not different between unprocessed and PEF processed orange juice (Elez-Martínez and others 2006). For grapes, a juice yield of 87%, similar to that after enzymatic maceration, and an increased content of soluble solids and pigments was reported after cell disintegration by PEF (Toepfl and others 2006). PEF treated cranberry juice had similar flavor and aroma profiles as the controls, and thermal treatment significantly altered the overall

flavor profile of the juice (Jin and Zhang 1999). No significant differences were observed in the content of anthocyanin pigments between PEF-treated samples and controls. However, thermal treatment significantly reduced the anthocyanin pigment content. HTST treatment of apple juice caused a considerable loss of phenols (32.2%) when compared with PEF treatment, which only caused a 14.49% reduction (Aguilar-Rosas and others 2007). The apple juice treated by PEF retained greater amounts of vitamin C and some representative flavor compounds, than the pasteurized juice (94.6°C for 30s) during storage at 4°C (Aguilar-Rosas and others 2007). Peroxidase and polyphenol oxidase showed a moderate 30-40% reduction with PEF treatment (Ho and others 1997). PEF treatment reduced 35% of PPO activity which was equivalent to heating the extract for 30 min at 60°C (Zuckerman and others 2004). Enzyme activity of PPO decreased 38.2% when treated at 33.6kV/cm for 126  $\mu$ s (Yang and others 2004)

### **High pressure processing**

HPP is considered a technology with the most promising perspective of industrial utilization. There are already known industrial applications, namely in Japan, USA, France and Spain (Houska and others 2006). The most successful products that are treated with high pressure pasteurization process are Golden oysters, orange juice, avocado sauce Guacamole, stewed packed ham, cooked rice and cooked reice mixtures, marinated chicken meat, etc. (Houska and others 2006). One of the main advantages of this process is the almost instantaneous and isostatic pressure transmission to the product, independent of size, shape, and food composition

yielding highly homogenous products (Deliza and others 2005). HPP can enable processors to produce innovative foods with fresh-like, natural-like attributes and natural looking colors which are all aspects valued by consumers (Deliza and others 2005). Information on the benefits offered by high pressure technology presented on the juice package also had a positive contribution on purchase intention (Deliza and others 2005).

High hydrostatic pressure is a promising alternative to traditional thermal processing techniques in food preservation, but associated changes to a diversity of phytonutrients have not been extensively investigated (Talcott and others 2003). Making comparisons between HPP and thermal processes is one way to assess its benefits by monitoring destruction of heat labile compounds (Talcott and others 2003).

High pressure treatment is expected to be less detrimental than thermal processing to low molecular weight food compounds such as flavouring agents, pigments, vitamins, etc., as covalent bondings are not affected by pressure (Butz and others 2002). The pressure stability of antioxidants is of interest since they reduce the risk of free radical-related oxidative damage that is associated with a number of diseases (Butz and others 2002). When used as a pretreatment of juice before processing, High Pressure treatment resulted in a significant increase in anthocyanins (Knorr 2003). In general, HPP was more detrimental to anthocyanins, ascorbic acid, and color characteristic as compared to thermally pasteurized and control juices due to oxidase enzymes that were active during HPP processing of muscadine juice (Talcott and others 2003). Potential mechanisms for destruction include the role of

PPO and/or autoxidative mechanisms, resulting in co-oxidation of anthocyanins and ascorbic acid (Talcott 2003). Without added ascorbic acid, total anthocyanin decreases were equal (3-5%) between treatments, but in the presence of ascorbic acid, anthocyanin losses were influenced by processing and resulted in 12.4% and 18.1% loss during pasteurization and HPP respectively (Talcott and others 2003). Losses ranged from 12 -15% following thermal pasteurization for delphinidin, cyanidin, petunidin and pelargonidin, compared to 15-25% for HPP while peonidin and malvidin were more stable (Talcott and others 2003). For high pressure to be a viable option for fresh muscadine grape juice, issues surrounding removal, inactivation or inhibition of native oxidase are critical for quality retention (Talcott and others 2003). Treatments of 400 MPa of muscadine juice had greater phytonutrients and antioxidant losses than treatment of 550 MPa due to the highly oxidative conditions that resulted from PPO activation during pressurization (Del Pozo-Insfran and others 2007). Anthocyanin degradation was observed at both processing pressures (400 and 550MPa) but was appreciably higher at 400MPa (70% loss) when compared with 550 MPa (46% loss), and these were correlated to antioxidant activity ( $r=0.86$ ) (Del Pozo-Insfran 2007). After 60 min of 600MPa pressure treatment and extreme temperature processing (95°C for 60 min), no changes were observed in the total concentration of lycopene of B-carotene when compared to control (Butz and others 2002). Stability of pigments may be explained by a matrix effect: within tissues, the pigments are often compartmentalized and thus protected from adverse influences (Butz and others 2002). Processing pressures of 500 and 800 KPa did not significantly reduced vitamin C and carotenoid content of treated orange juice when compared to

unprocessed orange juice (Fernandez Garcia and others 2001). Water soluble antioxidant capacities of orange and carrot juices can be increased by thermal treatment and maintained by high pressure treatment (Indrawati and others 2004).

CHAPTER III  
MATERIALS AND METHODS

**EXPERIMENT I**

**Extraction of anthocyanins and phenolics**

Frozen blueberries were homogenized in a commercial blender, weighed (25g), mixed with 25 ml of cold acetone (Fisher Scientific Co., Fair Lawn, NJ) and homogenized using a Brinkmann homogenizer (Polytron, Switzerland) at speed 4 for 2 minutes. An additional 25 ml of 70% acetone was used to wash the homogenizer and for reextraction. The extract was separated from insoluble plant material by filtering the slurry through a Whatman grade No.1 filter paper by vacuum suction using a Buchner funnel. Plant material was reextracted three times with 25 ml of 70% acetone until a clear filtrate was obtained. The volume of the filtrate was recorded and transferred to a separatory funnel where it was mixed with 2 volumes of chloroform (Fisher Scientific Co., Fair Lawn, NJ). The funnel was turned upside a down few times to mix solvents and stored overnight at 4°C until the clear portion between the two phases was obtained. The aqueous phase (upper portion) that contained the desired extract was transferred to a 500 ml boiling flask. The residual acetone/chloroform was removed in a Brinkman Bushi rotovapor model EL 131 (Brinkman Instruments, Westbury NY) at 40°C under vacuum.

The remaining aqueous extract was diluted to a known volume (100ml) with acidified deionized distilled water.

### **Total anthocyanins**

Total anthocyanins were determined using the pH differential method. To measure absorbance, the extract was diluted with buffer. The buffers used were pH =1 and pH=4.5. Buffer pH= 1 was made by mixing 125 ml of 0.2N KCl (Fisher Scientific Co., Fair Lawn, NJ) with 385 ml of 0.2N HCl (Fisher Scientific Co.). Buffer pH=4.5 was made by mixing 400ml of 1M sodium acetate (136g/l)(Fisher Scientific Co.) with 240 ml of 1N HCl and 360 ml distilled water. The pH of the buffers was adjusted as required to obtain final pH values of 1 and 4.5. The order of dilution was such that the sample at pH 1 had an absorbance of less than 1, preferably in the range of 0.4 to 0.6. The dilution strength was the same for both pH 1 and pH 4.5 samples. From the obtained extract, in order to obtain an absorbance in this range, the sample had to be diluted 50X, so the dilution factor was 50. The absorbance of the diluted extract was measured by placing the sample in a semimicro disposable cuvette (Fisher Scientific Co.) and reading the absorbance using a Genesys 5 UV/VIS spectrophotometer 336008 Model (Spectronic Analytical Instruments, Leeds, UK) at  $\lambda= 510\text{nm}$  and  $\lambda= 700\text{nm}$  for both buffers. The maximum absorbance for the anthocyanins was at  $\lambda= 510\text{nm}$ . Turbidity or haze was corrected by measuring the absorbance at  $\lambda= 700 \text{ nm}$  and subtracting this from the absorbance at the wavelength of maximum adsorption,  $\lambda= 510$ . Actual absorbance for the calculation of total anthocyanins was calculated using the following formula:

$$A = (A_{510} - A_{700})_{\text{pH } 1.0} - (A_{510} - A_{700})_{\text{pH } 4.5}$$

Concentration of total anthocyanins in mg/l was calculated using the following formula:

$$C \text{ (mg/l)} = A/\epsilon L \times \text{MW} \times 10^3 \times 50$$

where:

A = calculated absorbance

MW = molecular weight of cyanidin-3-glucoside (445,2 g/mol)

E = molar extinction coefficient of cyanidin-3-glucoside (26,900 M<sup>-1</sup> cm<sup>-1</sup>)

L = pathlength (1 cm)

### **Anthocyanin determination by HPLC-DAD and LC-MS**

An Agilent 1100 series HPLC (Agilent Technologies, Santa Clara CA) that is equipped with a quaternary pump, diode array detector, and a Gemini 5µm C18 110A (250 x 4.6 mm) column, fitted with a 4.0 x 3.0 mm inner diameter guard column, from Phenomenex (Torrance, Calif., U.S.A) was used for the separation of anthocyanins. Solvent A was 100% acetonitrile (Fisher Scientific Co.) and solvent B was 10% formic acid (Fisher Scientific Co.) in HPLC grade water (Fisher Scientific Co.). The program used the gradient presented in Table 1 with detection at 520 nm.

Table 1. Linear gradient program used for the separation of anthocyanins

Time (min)	Solvent %	
	Acetonitrile	10% formic acid
0	7	93
17	15	85
22	30	70
24	100	0
34	100	0

The injection volume was 25µl for pure extract of anthocyanin identification and 100 µl for diluted extract for anthocyanins degradation by PPO. The column was at room temperature and samples were filtered with 0.45 µm syringe filter (Fisher Scientific Co) before HPLC injection.

Anthocyanins were analyzed with LC-MS using Bruker Esquire(Bruker Daltonics, Bremen, Germany) in positive mode scanning from 100-500 m/z. located at Mississippi State Chemical Laboratory. The software employed was Esquire LC NT version 3.1 (Bruker Daltonics, Bremen, Germany). The program used same liner gradient like for HPLC separation (Table 1) except that solvent B was 0.1% formic acid in water.

### **Extraction of Polyphenoloxidase (PPO)**

Ten grams of frozen blueberries was mixed with 20 ml cold (~4°C) 200 mM phosphate buffer pH =7 containing 0.1% Triton 100-X (Sigma Chemical Co, St.Louis, MO) and 2% PVPP (Sigma Chemical Co.). The mixture was centrifuged at 10000 g for 15 min at 4°C using a Sorvall® RC 5B plus fixed angle centrifuge (Sorvall Products, L.P. Newton, CT). Triton 100-X was used to reverse or inhibit formation of tannin-protein complexes (Wesche-Ebeling and Montgomery 1990c). Blueberries contain a high level of phenolic compounds that may interfere in the extraction of the enzyme, or its enzymatic assay. Therefore PVPP, a not-specific phenolic absorbent, was included in the extraction buffer (Serradell and others 2000). The pellet was discarded and the supernatant was used for all further experiments as an enzyme source.

### Determination of total protein

Total protein was determined using the Micro-Bradford assay by placing 1 ml of 25 times diluted protein extract with 1 ml of Bradford reagent (Sigma Chemical Co.) directly in the cuvette. After 15 min of incubation, absorbance was recorded at 595 nm using a Genesys 5 UV/VIS spectrophotometer Model 336008 (Spectronic Analytical Instruments, Leeds, UK). Calculations of the total protein were made according to the standard curve (Figure 5). Standard curve was constructed by placing 1ml of a known concentration of BSA protein (Sigma Chemical Co.) with 1 ml of Bradford reagent and recording the absorbance at 595nm after 15 min of incubation.




Figure 5. Standard curve for determination of total protein with BSA as a standard

## **PPO assay**

Polyphenol oxidase activity was measured polarographically using a digital controller Model 10 equipped with a 7 ml Pyrex Rank inverted O<sub>2</sub> electrode (Rank Brothers, Cambridge, England) (Figure 6), and oxygen consumption was recorded using an ADC16 data logger (Pico Technology, St. Neots, UK). The instrument was standardized using air saturated water, assuming that 100% oxygen was equivalent to 233 μmol dissolved oxygen per liter (the solubility of O<sub>2</sub> in water equilibrated with air at 30°C) (Wrolstad 2001). To determine the electrode response and set zero oxygen concentration, few crystals of sodium dithionite (Sigma Chemical Co.) were placed in an electrode chamber that contained distilled water prior to performing the assay. In order to optimize the working concentration of the enzyme to ensure that assay gives a true measure of the initial rate of reaction, and that the relationship between enzyme concentration and rate lies within the linear range, the range finding was performed using different concentrations (amounts) of enzyme in the assay (Wrolstad 2001). The total assay volume was 3 ml containing 0.5ml (100mM) 4-methylcatechol (Sigma Chemical Co.), 0.2 ml enzyme extract and Mcillvane buffer pH=3.6. The temperature in the electrode incubation chamber was kept at 30°C by circulating water heated in the Isotemp 202 water bath (Fisher Scientific Co.) to 30°C with a pump (Manostat, New York, NY). The assay was performed for 180 s and the rate of reaction was calculated from the initial linear portion of the obtained curve. Results were expressed in μmol O<sub>2</sub>/min/100g of berries.




Figure 6. Digital controller model 10 (a) and Rank inverted oxygen electrode (b)

### **Inhibition study**

The polarographic method can be very useful in inhibition studies since it can indicate whether compounds were actually inhibiting PPO per se or merely affecting the formation of colored products (Ferrar and Walker 1996).

Food grade sodium benzoate, potassium sorbate (Xena International, St. Charles, IL), potassium metabisulfite (BASF Corporation, Florham Park, NJ) and a combination of sodium benzoate and potassium metabisulfite were used for the PPO inhibition study. Concentrations of inhibitors that were used are determined according to the standards currently used in the food industry (Table 2). Concentrations of 0.1% sodium benzoate and potassium sorbate are the maximum allowable concentrations (Title 21, U.S. Code of Federal Regulations ) for the use in food, while a concentration of  $\leq 10$  ppm of sulfite is not necessary for declaration on the label (Title 21, U.S. Code of Federal Regulations 101.100).

The effect of an inhibitor on PPO activity was determined by placing the designated amount of inhibitor solution in the electrode chamber together with 0.5ml (100mM) 4-methylcatechol (Sigma Chemical Co.), 0.2ml enzyme extract and McIlvane

buffer pH=3.6 to a total volume of 3 ml. Final concentrations of the inhibitors that were used are presented in Table 2. The temperature in the electrode incubation chamber was kept at 30°C by circulating water heated in the Isotemp 202 water bath (Fisher Scientific Co.) to 30°C with a pump (Manostat, New York, NY). The assay was performed for 180 s and the rate of reaction was calculated from the linear portion of the obtained curve. Activity of the enzyme was calculated in  $\mu\text{mol O}_2/\text{min}/100\text{g}$  of berries and expressed as a percentage compared to the enzyme activity without the inhibitor that was designated as 100%. Linear regression lines for each inhibitor were obtained by transforming percent activity to log percent activity and plotting that against inhibitor concentrations.

Table 2. Concentration of inhibitors used in the PPO inhibition study.

Inhibitors	Inhibitor levels				
	0	1	2	3	4
benzoate (%)	0	0.025	0.05	0.075	0.1
sorbate (%)	0	0.025	0.05	0.075	0.1
sulfite (ppm)	0	4	6	8	10
benzoate(%)/sulfite(ppm)	0	0.025/8	0.05/8	0.075/8	0.1/8

### Degradation of anthocyanins by blueberry PPO in the model system

The model system contained 400  $\mu\text{l}$  of anthocyanin extract ( $205.59 \pm 13.05$  mg/100g of blueberries); 200  $\mu\text{l}$  of crude enzyme extract ( $31.14 \pm 0.67$  mg/100g of blueberries of total protein and PPO activity of  $397 \pm 38$   $\mu\text{mol O}_2/\text{min}/100\text{g}$  of berries) and 500  $\mu\text{l}$  of 100mM 4-methyl catechol as a PPO substrate. Different volumes of inhibitors were added to make up final concentrations of 0.05 and 0.1 % of sodium benzoate and potassium sorbate, 4 and 10 ppm of potassium metabisulfite and 0.05 %/ 8ppm and 0.1%

/ 8ppm sodium benzoate/potassium metabisulfite. Total volume was adjusted to 3ml with Mcviline buffer pH 3.6. Total reaction time was 10 min at room temperature. After the reaction samples were filtered through a 0.45 micro filter (Fisher Scientific Co.) and analyzed with a HPLC utilizing the same linear gradient used for the separation and identification of anthocyanins (Table 1). Peak areas that were obtained with incubating just 400 µl of anthocyanin extract in a total volume of 3 ml in buffer was designated as 100 %. Anthocyanin extract was incubated separately for 10 min with the enzyme extract and 4-methyl catechol and no degradation of anthocyanin was observed.

### **Statistical analysis**

For the analysis of the inhibitor effect on the PPO activity, each inhibitor was treated as a separate experiment. Completely randomized designs with inhibitor concentrations as treatments with three replications were used to determine the effect of inhibitor concentration on PPO activity. The least significant difference test was used to separate treatments means ( $p < 0.05$ ). Slopes from the obtained regression lines for each inhibitor were treated as separate treatment and were analyzed in CR design and means were separated with LSD.

A split plot design was utilized for the analysis of anthocyanins destruction by PPO with and without the inhibitors. Inhibitor was designated as a main plot and the concentration of inhibitor as a split plot. Since the interaction between main and split plot was significant, mean separation was conducted on treatment combinations (inhibitor concentration) using SNK.

## **EXPERIMENT II**

### **Maceration pretreatment of blueberry mash**

Frozen blueberries were obtained from a local blueberry packer and were kept frozen (-15°C ) for about 10 months prior to processing. Blueberries were processed into juice at the Mississippi State University Dept. of Food Science Nutrition and Health Promotion Ammerman-Hearnsberger pilot plant. Frozen berries were heated in a steam kettle and constantly stirred during maceration. There were three maceration treatments (Figure 7) prior to pressing blueberries into juice: Control (no treatment), initial heat treatment and sodium benzoate treatment. Control fruit were thawed macerated at 55°C for 1h, pressed, placed into half gallon plastic jugs and frozen for further analysis. The heat treatment procedure differed only in that the fruit was heated to 90°C for 1 min prior to the maceration treatment. Sodium benzoate treatment differed from the control in that 0.1% of sodium benzoate (Xena International, St. Charles, IL) was added during the maceration step. All three treatments were replicated three times. Approximately 41 kg of frozen berries were used for each replication. After maceration, berries were pressed using a vertical press, and the juice was passed through a few layers of cheese cloth to remove seeds and pieces of skin. The juice was weighed to calculate the yield of the juice and then frozen for further treatment.

### **Blueberry juice filtration**

Before pasteurization, blueberry juice samples were thawed overnight in the refrigerator and first filtered through polypropylene string wound with a 100 micron

cartridge (FCPSF1100) and then through 20 micron (FCPSF1020) filters (Siemens Water Technologies, Lowell, MA) to remove particles. The resulting juice was frozen at -20°C until needed.

### **Juice treatments**

Frozen, filtered juice was defrosted overnight in a refrigerator prior to treatments. For heat pasteurization, thawed blueberry juice was processed in a steam jacketed kettle for 1 min at 90°C. Hot juice was immediately filled into 6 oz glass bottles and kept refrigerated (~3 d) at 4°C prior to analysis.

For Pulsed Electric treatment (PEF) thawed, blueberry juice samples, were transported to Louisiana State University in an ice chest (~ 5h) with ice and immediately processed. Pulsed Electric Field equipment was located at Louisiana State University in the Department of Dairy and Animal Science. Pulsed Electric field was carried out in a continuous flow bench scale system (OSU-4K, The Ohio State University, Columbus, OH) using square wave pulses (Figure 8). The flow rate was 60 ml/min and was controlled by a pump. The PEF processing conditions were 25 kV/cm electrical field applied in bipolar mode, 3  $\mu$ s pulse width, 10,000  $\mu$ s pulse period making a total of 600  $\mu$ s of pulse in 1 second ( $1000000 \text{ s} \times 6 \mu\text{s} / 10000 \text{ s}$ ) and 200 pulses per second (600  $\mu\text{s} / 3 \mu\text{s}$ ). Processed juice was filled in autoclaved 6 oz glass bottles. A portion of the filled bottled juice was frozen for further analysis.




Figure 7. Schematic representation of blueberry mash maceration treatments.

For High Pressure Processing (HPP), thawed blueberry juice (125ml) was vacuum packed in double bags and processed at 3 different pressures: 200, 400 and 600 MPa using a Quintus Food Press QFP 35-L600 with 7XS-6000 Intensifier Pump. These experiments were performed at Virginia Polytechnic Institute (Blacksburg, Virginia). The control sample was also vacuum packaged but was not subject to high pressure treatment. All samples were run with the water starting at room temperature (about 20°C) and with a holding time of 5 min. Pressures of 200, 400 and 600 MPa reached maximum vessel temperatures of 21, 27 and 32°C, respectively. Each pressure treatment was replicated three times.



Figure 8. OSU-4K Pulsed Electric Field Equipment.

### **Statistical analysis**

Completely randomized designs with three replications were utilized to analyze data obtained from maceration and juice processing treatments. The least significant difference test was used to separate treatments means ( $p < 0.05$ ).

### **Analysis**

#### **Extraction of anthocyanins and phenolics**

Frozen blueberry mash was homogenized in a commercial blender, weighed (25g) and processed as described in experiment I.

#### **Anthocyanin determination by HPLC-DAD**

Anthocyanins in the blueberry mash, pomace and juice were determined as described in experiment I. The only difference was that the concentration of anthocyanins was determined according to cyanidin-3-glucoside standard curve (Figure 9). An external standard was prepared by dissolving different amounts of cyanidin-3-glucoside in acidified HPLC grade water.




Figure 9. Cyanidin 3-glucoside standard curve

### **Extraction and assay of Polyphenoloxidase (PPO)**

Ten grams of frozen blueberry mash was used to determine PPO activity as described in experiment I.

### **Total phenolics**

Total phenolics in the blueberry extracts were determined with the Folin-Ciocalteu reagent by the method of Singleton and Rossi using gallic acid as the standard (Singleton and Rossi 1965). Twenty microliters of sample were mixed directly in the cuvette with 1.58 ml of water and 100  $\mu$ l of Folin-Ciocalteu reagent. The cuvette was incubated between 1 and 8 min and 300  $\mu$ l of 20% sodium carbonate was added. Samples were incubated at room temperature for 2 h and the absorbance was recorded at 765 nm using a Genesys 5 UV/VIS spectrophotometer Model 336008 (Spectronic

Analytical Instruments, Leeds, UK). Results were obtained using the gallic acid standard curve (Figure 10) and expressed as mg gallic acid equivalents in 100 g of blueberries.




Figure 10. Gallic acid standard curve.

### **Polymeric color**

Polymeric color was determined using the method described by Wrolstad (1976). The same extract that was used for the determination of anthocyanins and phenolics was diluted 50 times with distilled water. Two cuvettes were used for each sample. In each cuvette 2.8 ml of diluted sample were placed. In the first cuvette, 0.2 ml of freshly made 20% potassium metabisulfate (Fisher Scientific Co.) was added, and 0.2 ml of distilled water was placed in the second cuvette. Samples were left to equilibrate at room temperature (21°C) for 15 min. Absorbance for all samples was measured at  $\lambda=420\text{nm}$ , 510nm and 700nm (to correct for haze) against the blank cell filled with distilled water, using a Genesys 5 UV/VIS spectrophotometer Model 336008 (Spectronic Analytical

Instruments, Leeds, UK). Color density of the control sample (treated with water) was calculated using the formula;

$$CD = [ (A_{420} - A_{700}) - (A_{510} - A_{700}) ] \times 50$$

Polymeric color of the bisulfite bleached sample was calculated as follows:

$$PC = [A = (A_{510} - A_{700}) + (A_{510} - A_{700}) ] \times 50$$

Percent polymeric color was calculated as:

$$\% PC = (\text{Polymeric color} / \text{color density}) \times 100$$

### **Antioxidant activity**

The antioxidant activities of the extracts were determined using 2,2-Diphenyl-1-picrylhydrazyl (DPPH) (Sigma Chemical Co, St.Louis, MO) as a free radical, (Brand-Williams and others 1995). In its radical form, DPPH absorbs at 515 nm, but upon reduction by an antioxidant or radical species, the absorption disappears. Aliquots of 0.075 ml 5x diluted extract were added to 1.9 ml of 0.025g/l DPPH solution in methanol. The decrease in absorbance was determined at 515 nm at zero minutes, and every minute for a period of 20 min using a Genesys 5 UV/VIS spectrophotometer Model 336008 (Spectronic Analytical Instruments, Leeds, UK). The concentration of DPPH in the reaction medium was calculated from the calibration curve (Figure 11). The percentage of remaining DPPH (%DPPH<sub>REM</sub>) at the steady state, was determined as follows:

$$\%DPPH_{\text{REM}} = [DPPH]_T / [DPPH]_{T=0}$$

Results were expressed graphically by plotting time against the percentage of DPPH remaining after a given time.




Figure 11. 2,2-Diphenyl-1-picrylhydrazyl (DPPH) standard curve

## Acids

An Agilent 1100 series HPLC (Agilent Technologies, Santa Clara CA) equipped with quaternary pump, diode array detector and a Gemini 5 $\mu$ m C18 110A (250 x 4.6 mm) column, fitted with a 4.0 x 3.0 mm inner dia guard column, from Phenomenex (Torrance, Calif., U.S.A) was used for the separation of acids.

An isocratic elution with 0.01M H<sub>2</sub>SO<sub>4</sub> as the mobile phase, with the flow rate of 0.5 ml/min, column temperature of 35°C and UV detection at 214nm was used.

Standard curves of quinic (Figure 12), malic (Figure 13), citric (Figure 14) and shikimic acid (Figure 15) were used to calculate the concentration of these acids in the sample.




Figure 12. Quinic acid standard curve




Figure 13. Malic acid standard curve




Figure 14. Citric acid standard curve




Figure 15. Shikimic acid standard curve

## CHAPTER IV

### RESULTS AND DISCUSSION

#### **EXPERIMENT I**

##### **Enzyme activity and inhibition**

Significant differences were found between treatments (concentrations) for all inhibitors used ( $p < 0.01$ ). Addition of sulfite and combination of sulfite and benzoate had significantly higher regression slopes than benzoate alone and sorbate indicating better inhibition effect (Table 3).

Increasing the percentage of benzoate from 0 to 0.025 to 0.05 significantly decreased PPO activity (Figure 16). Utilization of 0.1 % sodium benzoate, which is the maximum that is allowed for use in food products (21CFR 184.1733), decreased PPO activity to only 5.05% in comparison to the activity without inhibitor (Figure 16). This PPO activity was not significantly different ( $p > 0.05$ ) from the activity of 8.78% when 0.075 % of benzoate was used.

When 4-methyl catechol is used as a substrate, benzoic acid is found to be a competitive inhibitor of PPO (Gunata and others 1987; Janovitz-Klapp and others 1990b). Benzoic acid at 2.5mM (0.03%) inhibited 43% of grape PPO at pH 5 (Gunata and others 1987) while in the present study a 76 % inhibition (23.98 % activity) was observed with 0.025 % at pH 3.6, using 4-methyl catechol as a substrate. The addition of

0.025% sodium benzoate reduced PPO activity to 23.98 % (Figure 16) while the same concentration of potassium sorbate reduced it to only 81.97 %, clearly indicating that the sodium benzoate is a better inhibitor. Benzoic and sorbic acid had a similar inhibition effect on PPO from banana, mushroom and apple, while sodium metabisulfite had the highest inhibition effect at 3300  $\mu$ M, the same concentration of (Ferrar and Walker 1996). The similarity in inhibition effect between benzoate and sorbate can be explained by the high concentrations used, 4.03% and 3.7% respectively, in comparison to 0.1% as the highest concentration used in our experiments. Potassium sorbate also significantly decreased PPO activity but with almost 50% of activity remaining when the maximum allowable concentration of 0.1% was used. Benzoic acid is a better inhibitor of gum arabic PPO compared to sorbic acid (Billaud and others 1996). Piffere and others (1974) also concluded that inhibitors containing the benzene nucleus showed a greater effectiveness than aliphatic compounds, with benzoic acid being a better inhibitor than sorbic acid. Of all the inhibitors used, potassium sorbate was the weakest PPO inhibitor with a significantly lower regression slope (Table 3). The presence of sorbic acid as an antimicrobial agent had a minimal effect on the browning of avocado puree (Soliva-Fortuny and others 2002).

Potassium metabisulfite in concentration as low as 4 ppm inhibited almost 70% of PPO activity (Figure 17). There was no significant difference ( $p > 0.05$ ) in PPO activity between 8 and 10 ppm sulfite (Figure 17), which is very important from the practical point of view since amounts of less than 10 ppm of sulfites do not have to be declared on the product label (Title 21, U.S. Code of Federal Regulations 101.100). Inhibition of PPO in the presence of sodium metabisulfite using an  $O_2$  -electrode indicated that the enzyme

was directly inhibited rather than by inhibition through secondary o-quinones. Ferrar and others (1996) concluded that sodium metabisulfite inhibition effect is probably due to the formation of colorless sulpho-quinones, rather than PPO inhibition per se, although the O<sub>2</sub> –electrode assay also showed inhibition, suggesting that enzyme inactivation may also occur. Direct and indirect effects of sulfites were reported by various authors and the mode of action depends on the inhibitor concentration, substrate and the complexity of the system (presence of other compounds).

Sodium metabisulfite was the most potent inhibitor of Emir grape PPO with 5.5, 13.5, 41.1 and 100% inhibition with concentrations of 0.05, 0.1, 0.25 and 0.50 mmol/L respectively (Unal and Sener 2006). Inhibition of PPO by sulfite in concentration of 0.04 mg/ml was instantaneous below pH 4 (Sayavedra-Soto and Montgomery 1986). Potassium metabisulfite at a concentration of 1mM inhibited 74% of strawberry PPO (Wesche-Ebeling and Montgomery 1990a). Metabisulfite at concentrations of 10 $\mu$ M, 0.1mM 0.4mM and 1mM inhibited 0, 31, 100 and 100% PPO activity from grapes respectively (Nunez-Delicado and others 2005). Na<sub>2</sub>S<sub>2</sub>O<sub>5</sub> inhibited 90, 98 and 100 % PPO activity from concord grapes in concentrations of 0.05, 0.5 and 5mM respectively (Cash and others 1976).

A combination of 0.05% of sodium benzoate with 8ppm of sulfite inhibited almost all PPO activity (Figure 18). There were significant differences between regression slopes between benzoate and the benzoate/sulfite combination, indicating that addition of 8ppm of sulfite had a significant effect on PPO inhibition (Table 3). From the present study it appears that the combination of benzoic acid and potassium metabisulfite (reducing agent) has a greater inhibition affect on blueberry PPO than either inhibitor by

itself. Similar results were obtained with the combination of benzoic acid and ascorbic acid (reducing agent) where browning of the apple was inhibited to a greater extent when a combination of inhibitors were used rather than either treatment alone. This demonstrates that the inhibition effect appears to be synergistic rather than additive (Sapers and others 1989). Ferrar and Walker (1996) also concluded that the majority of strong inhibitors were either aromatic or sulfhydryl compounds while aliphatic compounds exhibited weaker inhibition effect. This is in accordance to conclusions presented in this work since sodium benzoate and potassium metabisulfite were better inhibitors than potassium sorbate as demonstrated by the slopes of the inhibition (Table 3). Mechanism of inhibition may depend on the substrate and enzyme source (Ferrar and Walker 1996) which may explain some differences between results obtained in the current study and previous studies.

### **Identification of anthocyanins**

Anthocyanins were identified at 520 nm according to previous studies (Kader and others 1996; Lee and others 2002; Kahkonen and others 2003; Nakajima and others 2004; Buchert and others 2005). Peak identification was confirmed with HPLC/ESI-MS (Figure 20). Anthocyanins (mostly in the oxonium form at low pH) generate positive ions that can be detected by MS. An anthocyanin's single positive charge allows the mass-to-charge ratio ( $m/z$ ) to correspond directly to the molecular weight of the anthocyanins (Lee and Wrolstad 2004). Fifteen different anthocyanins were identified in the rabbiteye blueberry extract (Table 4) and the clear peaks can be seen on the extracted ion chromatogram (Figure 20). Fourteen anthocyanins were separated using

HPLC and had visible peaks at 520nm (Figure 19). Peonidin 3- arabinoside coeluted together with malvidin 3-glucoside (peak #13) and identification was only possible using mass spectroscopy. Lee and Wrolstad (2004) also found a low concentration of peonidin 3-arabinoside in highbush 'Rubel' blueberries which made its identification difficult. Coelution of the two can be clearly seen on the extracted ion chromatogram of anthocyanins, peaks 13 and 13a (Figure 20). In the same figure it can be seen that malvidin 3-galactoside and peonidin 3-glucoside also coelute, although their separation was possible using HPLC (Figure 19), possibly due to the use of 10% formic acid as opposed to only 0.1% formic acid used in HPLC-MS. Coelution of peonidin 3-glucoside with malvidin 3-galactoside (peaks 11 and 12) and peonidin 3-arabinoside and malvidin 3-glucoside (peaks 13 and 13a) was previously reported (Prior and others 2001; Lee and Wrolstad 2004; Nakajima and others 2004). Identical m/z and ion chromatograms of glycosides and galactosides of delphinidin (Figures 22 and 23), cyanidin (Figure 25 and 26), petunidin (Figures 31 and 32) peonidin (Figures 34 and 35) and malvidin (Figures 38 and 39) can be noted although they have different retention times. Their identification was possible due to the clear differences in retention times and the fact that galactosides elute before glycosides in the C18 column (Figures 21, 24, 30, 33 and 37). Extracted ion chromatograms and ion chromatograms of all 15 anthocyanins are presented in figures 21-41. Similar anthocyanin profiles to this study were reported for highbush blueberries grown in Europe (Kader and others 1996) and highbush "Rubel" blueberries grown in Oregon (Lee and Wrolstad 2004).

Derivatives of malvidin were the most abundant anthocyanins (42.1%), followed by delphinidin (19.86 %), cyanidin (11.58%), petunidin (16%) and peonidin (6.6 %).

Malvidin 3-galactoside was the most abundant individual anthocyanin, with 18.9%. Anthocyanins with the galactose in the molecule were the most abundant (44.68%) followed by the ones with the glucose (32.53%) and arabinose (18.93%).

Derivatives of malvidin and delphinidin accounted for 37 and 31% of the total anthocyanins, respectively, while 3-monogalactoside derivatives constituted about 41% (Kader and others 1996). Malvidin and delphinidin derivatives were the major anthocyanins in highbush blueberries (77.2%) (*Vaccinium corymbosum* L.cv. Rubel ) (Lee and others 2002). Malvidin 3-galactoside was a major anthocyanin in highbush ‘Rubel’ blueberries grown in Oregon with malvidin glycosides being most abundant accounting for 54.7% of all anthocyanins followed by delphinidin (25.7%), petunidin (14.7%), cyanidin (3.9%) and peonidin (1%) based on the peak area percentage (Lee and Wrolstad 2004). Malvidin 3-galactoside predominates in lowbush and ‘Tifblue’ blueberries (Prior and others 2001), in highbush ‘Rubel’ blueberries (Lee and others 2002) and rabbiteye blueberries grown in Japan (Nakajima and others 2004). Malvidin derivatives were most abundant in highbush blueberries, comprising 44% of the total anthocyanin in blueberries (Skrede and others 2000).

### **Degradation of anthocyanins**

In the model system, anthocyanins were degraded by the action of quinones formed by the action of blueberry PPO on 4-methyl catechol. Blueberry PPO degraded 36% of the total anthocyanins present in the model system (Figure 42). Degradation of anthocyanins was closely related to their structure. Triphenolic anthocyanins suffered the highest percentage of degradation (delphinidin 77 %), followed by diphenolic (petunidin

48% and cyanidin 24%) and monomeric (malvidin 19% and peonidin 16%) (Figure 43). All individual derivatives of these anthocyanins (glycosides, galactosides and arabinosides) followed a similar pattern of degradation (Figures 45-62).

The relative anthocyanin stability is related to their chemical structure. The delphinidin glycosides with the greatest liability have 3 ortho phenolic groups in the B ring and the cyanidin and petunidin derivative, which have the second order of reactivity, have 2 ortho phenolic groups. Peonidin and malvidin glycosides, which have the least reactivity, possess one phenolic substituent in the B ring with one and two adjacent methoxyl substituents, respectively (Skrede and others 2000). The high degradation of delphinidin anthocyanins can also be explained by the possible direct action of PPO. It was previously shown that PPO can act on anthocyanins when triphenolic function is present on the B ring of the flavylium structure as in the case of delphinidin (Sakamura and Obata 1963).

Fifty three percent of nasunin (delphinidin 3-(p-coumaroylrutinoside)-5-glucoside) was destroyed by PPO and the destruction increased to 98 % with the addition of chlorogenic acid (Sakamura and Obata 1963). This suggests that high degradation of delphinidin derivatives may be the result of direct action by PPO and indirect action by PPO generated quinones. Delphinidin glycosides were the most unstable during processing of the blueberry juice which mostly contributed to the action of blueberry PPO (Skrede and others 2000). Delphinidin-glycosides (based on its structure, delphinidin is most labile) were degraded the most, and malvidin-glycosides appear to have degraded the least by PPO, especially when the starting material was whole berries (Lee and Wrolstad 2004), which implies that the native enzymes present in the blueberry

destroyed anthocyanins (Skrede and others 2000). In a study of the Oxygen Radical Absorbance Capacity (ORAC) of different anthocyanins, it was reported that delphinidin and cyanidin derivatives had higher Trolox equivalents than malvidin and peonidin (Wang and others 1997), which is similar to results that were observed in this study. Wesche-Ebeling and Montgomery (1990) reported increased degradation of o-diphenolic anthocyanins in comparison to non o-diphenolic in a model system containing PPO and catechin as the PPO substrate. After 24 h, 60% of cyanidin anthocyanins was destroyed when compared to 50% pelargonidin anthocyanins (Wesche-Ebeling and Montgomery 1990b). In the presence of chlorogenic acid and enzyme extract, cyanidin-3- rutinoside was rapidly degraded and could not be detected after 20 min of reaction (Raynal and Moutounet 1989).

Malvidin derivatives should not be oxidized by the enzyme or by coupled oxidation (Sarni and others 1995). Degradation may occur due to the incorporation of anthocyanins into condensation products by quinone-phenol reactions as postulated by Wesche-Ebeling and Montgomery (1990). Sarni and others (1995) also demonstrated the degradation of both cyanidin and malvidin derivatives, with cyanidin degrading more rapidly than malvidin. The faster degradation of cyanidin-3-glucoside was due to its particular capacity (related to the o-diphenolic moiety) to be invoked in coupled oxidation, although the presence of methoxyl groups like in malvidin may also impede nucleophilic addition into quinone (Sarni and others 1995). Competition between cyanidin and malvidin 3-glucoside for the PPO generated o-quinone resulted in favor of the o-diphenol (cyanidin) (Sarni and others 1995), that can also explain more degradation of cyanidin in this study when compared to malvidin. Anthocyanins with arabinose in

their molecule seem to degrade more than the anthocyanins with glucose and galactose (Figure 44).

Addition of 0.05 and 0.1% sodium benzoate significantly decreased the percent of anthocyanins that were degraded by the PPO from 35% to 25% when 0.05% was used, and to 18 with 0.1% benzoate (Figure 42). With the addition of 0.1% sodium benzoate, almost 50% of degraded anthocyanins were protected. Protection of anthocyanin degradation by benzoate was not in the same order for all anthocyanins. The more the anthocyanins are susceptible to degradation the more they were protected by benzoate. At 0.1 %, benzoate protected approximately 30% of delphinidin (Figures 45-48), 25% of petunidin (Figures 49-52), 12% of cyanidin (Figures 53-56), 13% of malvidin (Figures 57-60) and 8% of peonidin (Figures 61-63). This is especially important since delphinidin and petunidin anthocyanins have the highest antioxidant activity and give the characteristic blue color to the blueberries.

Anthocyanin extracts contained additional compounds including acids and other phenolics. These compounds did not interfere with the inhibition of blueberry PPO by sodium benzoate. Even with an excess PPO substrate present, the generation of quinones that degrade anthocyanins is very fast, and benzoate was able to protect anthocyanins.

Addition of 8 ppm of potassium metabisulfite to the different concentrations of sodium benzoate (0.05% and 0.1%) did not significantly affect anthocyanin degradation compared to the addition of benzoate alone. Potassium metabisulfite and potassium sorbate did not have any protective effect on total and individual anthocyanin degradation by PPO. Potassium sorbate in the concentration of 0.1% significantly increased degradation of delphinidin (Figures 45-48) and petunidin (Figures 49-52) anthocyanins.

It is well known that sulfites can bleach anthocyanins, but since the order of degradation with sulfites was the same as without them, it can be concluded that the degradation of anthocyanins in the system with added sulfites was not due to the bleaching of anthocyanins but due to the action of PPO.

Although sulfites inhibited PPO activity (Figure 17), they did not protect anthocyanins that were present in the model solution. Concentration of sulfites (strong reducing agents) in our model system, might not be enough to reduce 4-methyl catechol quinones back to 4-methyl catechol which would protect anthocyanins. In the presence of excess reducing agent, o-quinines that were formed from the chlorogenic acid were reduced to o-diphenol (Kader and others 1999). Sulfur dioxide (8ppm) completely inhibited the degradation of anthocyanin in a model system containing mushroom PPO and catechol as a substrate at pH 6.5 (Goodman and Markakis 1965). In tart cherry juice, under similar conditions, but with no catechol added, 30ppm of SO<sub>2</sub> were required for a complete inhibition of anthocyanin degradation (Goodman and Markakis 1965). This greater concentration of SO<sub>2</sub> in the juice than in the model system necessary for the inhibition of PPO was attributed to the SO<sub>2</sub>-binding capacity of carbonyl compounds present in the juice. Influence of the added SO<sub>2</sub> on the content of the colored anthocyanins depends on the content of the SO<sub>2</sub>-binding carbonyl compounds present in wine with the same level of total SO<sub>2</sub> influencing differently anthocyanins color in different wines (Dallas and Laureano 1994). It was found that the amount of anthocyanins decreased less rapidly with the high concentration of SO<sub>2</sub> (Ribereau-Gayon and others 1983). Anthocyanin extract used in our model system contained other phenolic compounds and acids from the blueberries that might react with SO<sub>2</sub> before it

was able to inhibit PPO activity and formation of o-quinones that are responsible for anthocyanin destruction. Higher concentrations of potassium metabisulfite might be needed for the inhibition of anthocyanin degradation.

One of the reasons that sulfite did not inhibit the degradation of anthocyanins is that it may be in the bound form. Bound sulfite usually represents the portion of the additive which is present as hydroxysulphonate adducts that are formed by reactions between carbonyl groups and  $\text{HSO}_3^-$ . (Wedzicha 1992). Such adducts are decomposed slowly when sulfite treated foods are acidified or during titration with iodine, but rapidly if the pH of samples is raised to  $\text{pH} > 10$  or when acidified solutions are boiled (Wedzicha 1992). A well known observation of the browning of sulfite treated foods, while there is still residual sulfite present, indicates that in such situations the concentration of free sulphite may be depleted to an extent that it becomes kinetically limiting (Wedzicha 1992). The inhibition of enzymatic browning catalyzed by polyphenol oxidase is potentially capable of leading to irreversible binding of sulphites (Wedzicha 1992). In the case of 4-methyl catechol as a PPO substrate, sulfite can react with the formed o-quinone by forming 4-sulphocatechol which is neither oxidized by an enzyme nor does inhibit the enzyme (Wedzicha 1992).

Failure of potassium metabisulfite to protect against the degradation of anthocyanins could also be due to the fact that some of the sulfite was tight up in the enzyme sulfite complex after the enzyme had enough time to react with 4-methyl catechol and form quinones that reacted with anthocyanins.

Table 3. Slopes of the inhibitor regression lines.

Treatment	slope
Benzoate	0.307 b
Sorbate	0.078 c
Sulfite	0.54 a
Benzoate/sulfite	0.528 a
LSD	0.0785
CV	11.45

Means within the column followed by the same letter (abc) are not significantly Different ( $p < 0.05$ )




Figure 16. Effect of sodium benzoate and potassium sorbate on blueberry PPO activity

Means followed by the same letter (abcd and ABCD) are not significantly different ( $p < 0.05$ ).




Figure 17. Effect of potassium metabisulfite on blueberry PPO activity

Means followed by the same letter (abcd) are not significantly different ( $p < 0.05$ )




Figure 18. Effect of combination of sodium benzoate and potassium metabisulfite on blueberry PPO activity

Means followed by the same letter (abcd) are not significantly different ( $p < 0.05$ ).




Figure 19. HPLC chromatogram of 14 anthocyanins detected at 520 nm using DAD




Figure 20. Extracted ion chromatogram of 15 identified anthocyanins

Table 4. Retention times, % area and m/z of total and aglucone of identified anthocyanins in blueberry extract using High Pressure Liquid Chromatography

peak #	Name	Abbreviation	RT	% area	m/z total	m/z aglucone
1	Delphinidin-3-galactoside	Dp-3-gal	7.3	10	465	303
2	Delphinidin-3-glucoside	Dp-3-glc	8.24	4.83	465	303
3	Cyanidin-3-galactoside	Cy-3-gal	9.2	5.68	449	287
4	Delphinidin-3-arabinoside	Dp-3-ara	9.62	5.03	435	303
5	Cyanidin-3-glucoside	Cy-3-glc	10.5	3.3	449	287
6	Petunidin-3-galactoside	Pt-3-gal	11.2	7.3	479	317
7	Cyanidin-3-arabinoside	Cy-3-ara	11.83	2.6	419	287
8	Petunidin-3-glucoside	Pt-3-glc	12.4	5.6	479	317
9	Peonidin-3-galactoside	Pn-3-gal	13.5	2.8	463	301
10	Petunidin-3-arabinoside	Pt-3--ara	14.02	3.1	449	317
11	Peonidin-3-glucoside	Pn-3-glc	15.03	3.8	463	301
12	Malvidin-3-galactoside	Mv-3-gal	15.33	18.9	493	331
13	Malvidin-3-glucoside	Mv-3-glc	16.7	15	493	331
13a*	Peonidin-3-arabinoside*	Pn-3-ara	16.7		433	301
14	Malvidin-3-arabinoside	Mv-3-ara	18.4	8.2	463	331




Figure 21. Extracted chromatogram of delphinidin 3-galactoside and delphinidin 3-glucoside m/z 465




Figure 22. Ion chromatogram of delphinidin 3-galactoside (peak # 1)




Figure 23. Ion chromatogram of delphinidin 3-glucoside (peak # 2)




Figure 24. Extracted chromatogram of cyanidin 3-galactoside, cyanidin 3-glucoside and petunidin 3-arabinoside m/z 449




Figure 25. Ion chromatogram of cyanidin 3-galactoside (peak # 3)




Figure 26. Ion chromatogram of cyanidin 3-glucoside (peak # 5)




Figure 27 . Ion chromatogram of petunidin 3-arabioside (peak # 10).




Figure 28. Extracted chromatogram of delphinidin 3- arabioside m/z 435




Figure 29. Ion chromatogram of delphinidin 3- arabioside (peak # 4)




Figure 30. Extracted chromatogram of petunidin 3-galactoside and petunidin 3-glucoside m/z 479.




Figure 31. Ion chromatogram of petunidin 3-galactoside (peak # 6)




Figure 32. Ion chromatogram of petunidin 3-glucoside (peak # 8)




Figure 33. Extracted chromatogram of peonidin 3-galactoside, peonidin 3-glucoside and malvidin 3-arabinoside m/z 463




Figure 34. Ion chromatogram of peonidin 3-galactoside (peak # 9)




Figure 35. Ion chromatogram of peonidin 3-glucoside (peak # 11)




Figure 36. Ion chromatogram of malvidin 3-arabinoside (peak # 14)




Figure 37. Extracted chromatograms of malvidin 3-galactoside and malvidin 3-glucoside m/z 493




Figure 38. Ion chromatogram of malvidin 3-galactoside (peak #12)




Figure 39. Ion chromatogram of malvidin 3-glucoside (peak #13)




Figure 40. Extracted chromatogram of peonidin 3-arabinoside m/z 433




Figure 41. Ion chromatogram of peonidin 3-arabinoside




Figure 42. Degradation of total anthocyanins by blueberry PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 43. Degradation of anthocyanins by blueberry PPO




Figure 44. Degradation of anthocyanins by PPO grouped according to the sugar in the molecule




Figure 45. Degradation of delphinidin anthocyanins by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 46. Degradation of delphinidin 3-galactoside by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 47. Degradation of delphinidin 3-glucoside by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 48. Degradation of delphinidin 3-arabinoxide by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 49. Degradation of petunidin anthocyanins by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 50. Degradation of petunidin 3-galactoside by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 51. Degradation of petunidin 3-glucoside by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 52. Degradation of petunidin 3-arabinoside by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 53. Degradation of cyanidin anthocyanins by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 54. Degradation of cyanidin 3-galactoside by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 55. Degradation of cyanidin 3-glucoside by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 56. Degradation of cyanidin 3-arabinoxide by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 57. Degradation of malvidin anthocyanins by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 58. Degradation of malvidin 3-galactoside by PPO with and without Inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 59. Degradation of malvidin 3-glucoside by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 60. Degradation of malvidin 3-arabinoxide by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 61. Degradation of peonidin anthocyanins by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 62. Degradation of peonidin 3-galactoside by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 63. Degradation of peonidin 3-glucoside by PPO with and without inhibitors

Means followed by the same letter are not significantly different ( $p < 0.05$ )

## EXPERIMENT II

### Identification and quantification of anthocyanins

Fourteen different anthocyanins were identified in the frozen rabbitey blueberry extract (Table 4) and clear peak separations can be seen on the HPLC chromatogram detected at 520 nm (Figure 19). Separation of Peonidin 3- glucoside and malvidin 3- galactoside, peaks number 11 and 12 (Figure 19) was variable. In most of the analysis for different treatments, the two coeluted and no clear pattern of separation could be found. Because of that, it was impossible to analyze data for peonidin 3- glucoside, so the data was omitted in all analysis. Determination of total anthocyanins by HPLC method was a summation of individual anthocyanin peaks detected at 520 nm. Individual anthocyanins were quantified by comparisons with an external standard of cyanidin-3- glucoside.

Concentration of total anthocyanins in frozen blueberries was 328 mg/100 g. These value fall within the range of 25 to 495mg/100g for highbush blueberries (Mazza and Miniati 1993). In the frozen berries, malvidin glucoside was the most abundant (50.43%) followed by delphinidin (19.3), petunidin (15.64), cyanidin (10.68%) and peonidin (4.1%). Concentrations of individual anthocyanins expressed as mg/L of cyanidin 3–glucoside are presented in table 3. Malvidin and delphinidin derivatives were the major anthocyanins in highbush blueberries (77.2%) (*Vaccinium corymbosum* L.cv. Rubel ) (Lee and others 2002). Malvidin 3-galactoside was a major anthocyanin in highbush 'Rubel' blueberries grown in Oregon with malvidin glycosides being most abundant accounting for 54.7% of all anthocyanins followed by delphinidin (25.7%),

petunidin (14.7%), cyanidin (3.9%) and peonidin (1%) based on the peak area percentage (Lee and Wrolstad 2004). Malvidin 3-galactoside predominates in lowbush (*V. angustifolium*) and ‘Tifblue’ (*V. ashei*) blueberries (Prior and others 2001), in Highbush ‘Rubel’ blueberries (Lee and others 2002) and rabbiteye (*V. ashei*) blueberries grown in Japan (Nakajima and others 2004). Malvidin derivatives were most abundant in highbush blueberries, comprising 44% of the total anthocyanins in blueberries (Skrede and others 2000).

### **Identification and quantification of acids**

Separation of acid standards (Figure 64) and acids in the blueberry extract (Figure 65) was carried out at 214 nm. The acids that were identified in frozen blueberry extract were quinic, malic, citric and shikimic. The most abundant acid identified in the blueberry extract was quinic 941.6 mg/100g, followed by malic 89.4 mg/100g, citric 29.4 mg/100g and shikimic 8.9mg/100g (Table 6). Conflicting reports about acid concentrations were found in the literature. In rabbiteye fruit, the percentage contribution of citric, succinic, malic and quinic acid is about 10%, 50%, 34% and 6%, respectively (Kalt and others 1996). In the present research, blueberry extract was also screened for succinic acid but it was not detected. Other researchers reported an average of 95% citric acid and 1% to 2% each of quinic and malic acid in ripe “Wolcott” fruit (Kushman and Ballinger 1968). Succinic and malic acids were the predominant acids in rabbiteye cultivars averaging 50% and 33%, respectively (Ehlenfeldt and others 1994). Among rabbiteye clones, citric acid averaged 10% and was never found to be more than 22%,

while quinic acid was consistently present only as a minor constituent, averaging 6% (Ehlenfeldt and others 1994).

### **PPO activity in blueberry mash**

Maceration treatment affected ( $p < 0.05$ ) PPO activity (Figure 70). Maceration temperature at 55°C reduced ( $p < 0.05$ ) PPO activity from 233 to 52.3  $\mu\text{molO}_2/\text{min}/100\text{g}$ . Although there were no significant differences between the control treatment (60 min maceration at 55°C) and heat treatment, no PPO activity was detected in blueberry mash that was heat treated at 90°C and then macerated for 60 min at 55°C. This is to be expected since PPO is not an extremely heat-stable enzyme. Short exposures, in the tissue and solutions, to temperatures of 70 to 90°C are, in most cases, sufficient for partial or total irreversible destruction of its catalytic function (Vamos-Vigyazo 1981). Blanching blueberry fruit for 3 min in a steam tunnel totally inactivated PPO (Rossi and others 2003). Inactivation of PPO did not begin to occur until the grape juice was heated above 60°C and heating above 87°C was necessary to completely inactivate the enzyme (Montgomery and others 1982). Activity of the PPO was reduced 100 times when the temperature was raised to 88-99°C, to 4 and 3  $\mu\text{moles O}_2/\text{min}/2.5\text{ml}$  from 445  $\mu\text{moles O}_2/\text{min}/2.5\text{ml}$ , while at 60°C PPO activity decreased 10 times to 44  $\mu\text{moles O}_2/\text{min}/2.5\text{ml}$  (Montgomery and others 1982). The PPO in crude extracts of plum, apple, pear and avocado was inactivated at 60-65°C, whereas in grape extracts, 55°C was enough for inactivation (Weemaes and others 1998).

Addition of 0.1% of sodium benzoate to blueberry mash significantly reduced PPO activity compared to frozen blueberries but was also unexpectedly significantly

higher than a control sample without added inhibitor (Figure 70). It could be possible that the addition of sodium benzoate increased thermal stability of PPO and that the added amounts were not enough to inhibit enzyme. Thermal stability of PPO can be affected by some substances present in the food (Tomas-Barberan and Espin 2001a). Addition of both EDTA and benzoic acid increased the thermal stability of mushroom PPO, whereas glutathione produced a sensitization to temperature treatments, probably due to an interaction with a disulphide bond in the enzyme (Weemaes and others 1997). Benzoic acid at 50 mM protected PPO from temperature treatment, thus increasing the thermal stability of the enzyme (Weemaes and others 1997). It could also be possible that some other phenolic compounds that are abundant in blueberries are bound to PPO without inhibiting activity but preventing sodium benzoate to bind to an enzyme and inhibit its activity. There are two distinct substrate binding sites on the enzyme molecule, one of which had a high affinity for aromatic compounds, including phenolic substrates, while the other, which probably contained the enzyme copper, served for metal-binding agents and oxygen (Duckworth and Coleman 1970). In the system with benzoic acid, phenolic substrate (4-methyl catechol) and oxygen are present, oxygen is the first substrate to be bound to the apple PPO forming E-O<sub>2</sub> complex to which benzoic acid as an inhibitor can bind only with enzymatic forms of PPO which are free of other phenolics i.e., E or E-O<sub>2</sub> (Janovitz-Klapp and others 1990b).

### **Degradation of anthocyanins, acids and phenolics in blueberry mash**

Since blueberries contain anthocyanins that impart their color only in the skin, breaking the skin and maceration of the mash are necessary to extract the color and

obtain good quality juice. Freezing and heating of the blueberries had a more pronounced effect than pectolytic enzymes on the skin cells, and consequently on the liberation of color pigments. The loss of semipermeability and mechanical injury to the cell membranes appears to determine the total color released (Fuleki and Hope 1964). Significant differences were found between maceration treatments in respect to concentration of total anthocyanins (Figure 66). No significant differences were found between frozen blueberries and control mash treatment (Figure 66), proving that 55°C was not high enough to affect anthocyanins but on the other hand was high enough to decrease PPO activity almost 4 times (Figure 70), thus protecting anthocyanins and preserving the red color.

Blueberry puree that was held for 60 min at 60°C had higher concentrations of anthocyanins and phenolics than the one held at 25°C, with anthocyanins being more affected than phenolics (Kalt and others 2000). The increased extraction of anthocyanins was attributed to the increased permeability of membranes in the macerated berries at higher temperatures and to decreased solubility of oxygen at higher temperatures that can prevent oxidative degradation (Kalt and others 2000). After 2h mash treatment at 50°C, 97% of the original anthocyanin content in black currant berries were intact (Iversen 1999).

Maceration treatments with added sodium benzoate and heat had a significantly lower concentration of total anthocyanins (Figure 66). Heat treatment at 90°C inactivated PPO, but degraded approximately 30% of anthocyanins. The kinetics of thermal degradation is generally first order up to 110°C and is influenced by temperature, plant species, pH, presence of oxygen and sugar content. A logarithmic relationship between

temperature and pigment degradation has been described (Francis 1989). Although PPO activity can degrade anthocyanins, degradation of anthocyanins in the heat treated (blanched) blueberry mash can be the result of the heat treatment rather than the action of PPO since PPO activity was barely detectible in the heat treated mash (Figure 70).

The benzoate treatment also significantly reduced total anthocyanins, but possibly due to the action of PPO rather than heat, since 55°C was not enough to degrade anthocyanins (Figure 66). Addition of sodium benzoate decreased PPO activity only by 42% from 233 to 137  $\mu\text{molO}_2/\text{min}/100\text{g}$  (Figure 70) leaving enough active enzyme to degrade anthocyanins. It was shown that substantial losses of anthocyanins and other polyphenolics occurred when blueberries were processed into juice and that different classes of compounds had varying susceptibility to degradation with different processing operations with the highest losses occurring during milling and depectinization due to the action of native PPO (Skrede and others 2000).

Not all anthocyanin aglucones were effected the same by the treatments. No significant differences were found between frozen blueberries and control mash in respect to all anthocyanin aglucones (Figure 67). All anthocyanin aglucones except cyanidin aglucones were significantly decreased by heat treatment (Figure 67) when compared to frozen berries. Heat treatment had the highest negative influence on malvidin anthocyanins (~34 % degraded), followed by delphinidin and petunidin (28% degraded), and cyanidin (20% degraded), in comparison to the frozen berries (Figure 67). The three most representative blueberry anthocyanins (i.e. malvidin glycosides) showed a minor recovery increase with blanching (Rossi and others 2003). Sodium benzoate significantly decreased all anthocyanin aglucones due to the action of PPO that was still

active (Figure 67). Treatment with benzoate showed a similar trend of anthocyanin degradation except that malvidin glycosides were less degraded than with the heat treatment, with only 25% degradation. These results suggest that malvidin glycosides are more susceptible to heat degradation than to the action of PPO. Malvidin glycosides are non o-diphenolic anthocyanins which are less susceptible to degradation of PPO generated o-quinines. Nevertheless, Sarni and others (1995) noticed a decrease in malvidin-3-glucoside in a model system in the presence of PPO and caffeoyltartaric acid (PPO substrate).

Individual anthocyanins were affected by maceration treatment similarly to their aglucones (Table 5). Maceration treatments did not have a significant influence on polymeric color (Figure 69). This indicates that the association of anthocyanins with each other and with other phenolics occurred within the blueberries and was not affected in any way by treatments.

No significant differences were found between maceration treatments in respect to total phenolic concentration (Figure 68). Total phenolic concentration was retained during jam processing of some berries (Amakura and others 2000; Zafrilla and others 2001). Thermal processing of raspberries into jam had a minimal effect on the major phenolics present (Zafrilla and others 2001).

Significant differences were found between blueberry mash maceration treatments in respect to the concentration of identified acids (Table 6). Concentrations of quinic acid and citric acids were significantly higher in heat treated mash than in the control maceration treatment (Table 6). Heat treatment probably helped increase the extraction of these acids from the blueberry skin and seeds.

## **Juice yield**

Yield of blueberry juice was around 52% and there were no significant differences between maceration treatments (Figure 75). Higher yields for blueberry juice were reported in previous studies: 79-81 % (Rossi and others 2003), 83% (Roberts and others 2004), 83%(Skrede and others 2000), and 78-82% for both cold and hot pressed juice (Carlson 2003). Higher yields may be due to the use of added pectolytic enzymes, smaller scale production and use of more sophisticated pressing equipment.

## **Anthocyanins, phenolics and acids in pressed juice and press cake**

After pressing, anthocyanins are released into juice either in soluble form or in small suspended fruit particles (Iversen 1999). No significant differences in anthocyanins were found between different maceration treatments in extracted juice (Figure 71). Although, heat treated blueberry mash had a significantly lower concentration of total anthocyanins after 60 min of maceration when compared to the control (Figure 66), no significant differences in anthocyanins in the extracted juice could be the consequence of better extraction of anthocyanins from the heat-broken skin. Recovery of anthocyanins was 12 % from the control and benzoate treated mash, and 17% from the heat treated juice. Anthocyanins are not as efficiently extracted in the pressing operation as are sugars, acids and other water solubles, which can have a negative impact on juice quality (Skrede and others 2000; Lee and others 2002). It was shown that substantial losses of anthocyanins and other polyphenolics occurred when blueberries were processed into juice and that different classes of compounds had varying susceptibility to degradation

with different processing operations, with the highest losses occurring during milling and depectinization due to the action of native PPO (Skrede and others 2000).

Pretreatments with heat and SO<sub>2</sub> resulted in higher recovery of red color pigments with a more intense color (higher chroma) in pasteurized juice when compared to the control, while the heat treated samples were darker (smaller L values) than the control and SO<sub>2</sub> treated juice (Lee and others 2002). Only 12% of total blueberry anthocyanins were extracted in juice, increasing to 23% when the juice was blanched (Rossi and others 2003). Rossi and others (2003) reported that the total anthocyanin content of juice from blanched blueberry fruit was twice that of non blanched fruit. These researchers concluded that this was the result of PPO inactivation and the greater extraction yield was linked to the increase of fruit skin permeability caused by the heat treatment.

No significant differences in anthocyanin aglucones were found between control extracted, benzoate and heat treated juices (Figure 72). Heat extracted juice had significantly higher amounts of delphinidin, cyanidin and petunidin anthocyanins compared to benzoate treated juice (Figure 72). Malvidin glycosides were the most readily extracted anthocyanins for all treatments, contributing approximately 57-65% of the total anthocyanins that were present in the juice, while their contribution in the frozen blueberry fruit was approximately 50%. Malvidin glycosides comprised 44% of the total anthocyanins in blueberry fruit and increased to 63% in pasteurized juice and concentrate (Skrede and others 2000). Similar results were reported by Lee and others (2002) where malvidin glycosides in the berry were 51% of total anthocyanins and increased to 60-77% in pasteurized juice and concentrate. Contribution of delphinidin and petunidin anthocyanins to total anthocyanins decreased in extracted juice compared to their

contribution in frozen berries, while cyanidin and peonidin ratios remained the same. Delphinidin anthocyanins decreased the most of all anthocyanins, decreasing from 19% contribution of total anthocyanins present in the frozen berries to 11% in the pressed juice. Delphinidin glucosides were the most unstable decreasing from 12% in blueberry fruit to 5% in pasteurized juice with only trace amounts in the initial pressed juice (Skrede and others 2000). Decreases in delphinidin glycosides was also reported by Lee and others (2002) but only for the juice that received no pretreatment before pressing. There was a concomitant decrease in delphinidin and petunidin glycosides (Lee and others 2002). The proportion of delphinidin-glycosides decreased especially in the control sample with only 8% left, while heat and SO<sub>2</sub> treated samples had 20 and 23% retained in the pasteurized juice (Lee and others 2002). The relative ratio of delphinidin glycosides versus cyanidin glycosides remained about the same in the juices as compared with the whole berry (Buchert and others 2005). Rossi and others (2003) obtained very different results with the higher percent recovery increase of delphinidin glycosides and petunidin –glycosides which have a more intense blue than that of cyanidin-glycosides and peonidin-glycosides, which are orange in color. The highest percentage recovery increase observed for delphinidin-glycosides could also be linked to their higher water-solubility, due to the presence of three free phenolic functions (Rossi and others 2003).

The anthocyanins that showed the highest benefits from the blanching treatment were, in decreasing order, the glycosides of delphinidin, petunidin , and cyanidin with the exception of cyanidin-3-glucoside that showed the lowest recovery (Rossi and others 2003). The three most representative blueberry anthocyanins (i.e. malvidin glycosides) showed a minor recovery increase with blanching (Rossi and others 2003). In the present

study, heat treatment (blanching) did not significantly decrease recovery of malvidin glycosides (Figure 67). Lee and others (2002) concluded that heat and SO<sub>2</sub> treatment of highbush blueberries were effective in increasing the recovery of anthocyanins but not other phenolics in the juice. Only significant differences between treatments in extracted juice with respect to individual anthocyanins were found between delphinidin-3-galactoside, cyanidin-3-galactoside, petunidin-3-galactoside and malvidin-3-arabinoside (Table 7). There were no significant differences in percent polymerization between different treatments indicating that anthocyanins did not associate with each other or with other phenolics (Figure 74).

Substantial amounts of anthocyanins, 641, 595 and 503 mg/100g of press cake remained in the control, benzoate and heat pretreated press cake after the extraction of juice (Figure 76). No significant differences in the press cake were found between different pretreatments in respect to total anthocyanins. Heat pretreated press cake had significantly lower concentration of malvidin glycosides (Figure 77). Concentration of malvidin -3- galactoside and malvidin-3-arabinoside was significantly lower in the heat pretreated press cake (Table 9). This low concentration was expected since malvidin glycosides were the most readily extracted anthocyanins for all treatments, contributing approximately 57-65% of total anthocyanins present in the juice, while their contribution in the frozen blueberry fruit was around 50%. In addition, the heat treatment had the highest negative influence on malvidin anthocyanins (~34 % degraded), followed by delphinidin and petunidin (28% degraded) and cyanidin (20% degraded) in comparison to the frozen berries (Figure 67). Changes in malvidin glycosides with processing were the most noticeable, ranging from 50% in frozen berries, to 65% in pressed juice and to 43%

in the press cake. The press cake residues contained substantial amounts of anthocyanins, about 43-55% of frozen berry anthocyanins, with the control press cake containing the greatest amounts (Lee and others 2002). No significant differences in percent anthocyanin polymerization in press cake were found between different pretreatments (Figure 79).

Significant differences were found between pretreatments in respect to the total amount of phenolics extracted into the juice (Figure 73). Heat extracted juice had a significantly higher amount of total phenolics than the benzoate treatment, but not more than the control (Figure 73). Less extraction of phenolics from the benzoate treated blueberry mash can be explained by the fact that benzoate treated mash had a lower initial concentration of phenolics (Figure 68). Phenolics percent extraction from the corresponding blueberry mash was 33%, 35% and 41% for the control, benzoate treated and heat treated samples, respectively. Lee and others (2002) reported a considerable loss in phenolics during thawing, crushing and pressing with similar results obtained by Skrede and others (2000). Sixty to 65% of blueberry polyphenolics were lost during thawing, crushing and pressing (Lee and others 2002). There were not significant differences in polyphenolics taken after each processing step between the control, heat and SO<sub>2</sub> treated berries. Thirty six to 39% of the polyphenolics in the berries were present in the pasteurized juice. Forty two to 45% of the frozen berry phenolics were lost during juice processing and not accounted in the final pasteurized juices and press cake (Lee and others 2002). Control pressed juice (no pretreatment) and SO<sub>2</sub> treated pressed juice had lower polyphenolics than their pasteurized juices. Hot pressed blueberries had higher amounts of total phenolics than the cold pressed ones (Lee and others 2002).

Considerable amounts of total phenolics remained in the press cake after juice extraction. Heat pretreated press cake seem to contain the highest amount of phenolics, but was not significantly higher than the other two treatments (Figure 78). Blueberry extract had 5500 ppm of total phenolics, hot pressed blueberries had around 3200 ppm while pasteurized blueberries had 5000 ppm (Carlson 2003).

The blueberry mash treatment did not have a significant influence on extracted acids, except for citric acid (Table 8). Heat pretreated extracted juice had significantly higher amounts of citric acid when compared to control and benzoate pretreated juice (Table 8). The higher amounts of extracted citric acid are the result of higher amounts of citric acid in the blueberry mash from which the juice was extracted (Table 6). Heat treatment probably broke the cells which helped extract more citric acid. The press cake did not contain any citric acid for any of the treatments, indicating that this acid is readily extracted into juice (Table 10). Significant differences between treatments were found with respect to concentration of shikimic and malic acid in the pomace (Table 10). Their concentrations were higher in the heat-treated pomace and lower in the benzoate-treated pomace.

## **Juice Processing**

### **Pulsed electric fields, PEF**

Many researchers indicate that PEF operating with electric fields between 15 and 80kV/cm, pulse widths between 1 and 100Ms, and a frequency of 1-100 pulses, were sufficient for inactivation of bacteria and certain enzymes, while taste and flavors were unaffected (Barbosa-Canovas and others 2001; Wesierska and Trziszka 2007). The PEF

treatment did not have significant influence on total anthocyanins, individual anthocyanins, total phenolics, percent anthocyanin polymerization, and DPPH remaining after 5 min (Table 11). In addition to that, no significant differences were found in respect to anthocyanins aglucones (Figure 80). It was observed that PEF-pretreatment caused an increase in anthocyanin concentration in grape juice (Knorr 2003). The PEF treatment caused degradation of cyanidin-3 glycoside in a aqueous-methanol solution. As the electric field intensity and the treatment time increased, degradation of cyanidin-3 glycoside significantly increased (Zhang and others 2007). Purified solution of single anthocyanins might be more susceptible to degradation than a juice extract containing other soluble compounds. PEF-treated cranberry juice had a similar flavor or aroma profiles as the controls, while thermal treatment significantly altered the overall flavor profile of the juice (Jin and Zhang 1999). No significant differences were observed in the content of anthocyanin pigments between PEF-treated samples and controls. However, thermal treatment significantly reduced the anthocyanin pigment content. HTST treatment of apple juice caused a considerable loss of phenols (32.2%) when compared with PEF treatment, which only caused a 14.49% reduction (Aguilar-Rosas and others 2007). Antioxidant activity monitored as percent DPPH over a period of 20 min remained the same before and after PEF treatment (Figure 81). Antioxidant activities that were determined with the DPPH method were not different between unprocessed and PEF processed orange juice (Elez-Martinez and others 2006). The PEF treatment also did not have any significant influence on the acids that are present in the juice (Figure 82). Organic acid and volatile aroma compounds of citrus juices did not change with PEF treatment at 28kV/cm with 50 pulses (Cserhalmi and others 2006).

## **Pasteurization**

Kettle pasteurization for 1 min at 90°C did not have a significant influence on anthocyanins aglucones (Figure 83), total anthocyanins, individual anthocyanins, % anthocyanin polymerization, total phenolic and % DPPH remaining after 5 min (Table 12). There are conflicting reports in the literature on the influence of heat on phytonutrients in red colored juices. Industrial processing of pomegranates to obtain juices increased their antioxidant capacity and phenolics content (Gil and others 2000). Reduction of anthocyanins content in strawberry juice upon heating was reported (Rwabahizi and Wrolstad 1988 and Bakker and others 1992). Juice obtained after filtration was not clear blueberry juice which could have some protective effect on phytonutrients present during pasteurization. Large loss of strawberry anthocyanins after clarification of the juice compared to the purees was observed, and it was attributed mostly to the loss of colored compounds bound to the proteins due to filtration (Bakker and others 1992). Pigment destruction was more pronounced in clarified juice (Wrolstad and others 1994).

Curves obtained by monitoring the disappearance of DPPH could not be distinguished, indicating the same antioxidant activity of juice before and after pasteurization (Figure 84). The DPPH free radical scavenging activity assay did not show clear division between heat-treated and non-heat treated products (Schmidt and others 2005).

Pasteurization using a steam kettle was the most effective at retaining antioxidant activity as measured by total phenols and (ORAC) in the final products. One likely reason for this significantly higher level of antioxidant retention is the exposure to higher

heating temperatures resulting in increased permeability of water soluble substances, moving out of the cells. The higher heat also degraded enzymes (PPO and glucosidase) present that would be harmful to the anthocyanins and also lowered the levels of oxygen present in the final products which contribute to antioxidant destruction (Carlson 2003). Pasteurization caused a significant increase of quinic and shikimic acids (Figure 85).

### **High Pressure Processing, HPP**

High pressure treatment is expected to be less detrimental than thermal process to low molecular weight food compounds such as flavouring agents, pigments, vitamins, etc., as covalent bondings are not affected by pressure ( Butz and others 2002). Although different processing pressures did not have a significant influence on concentration of total anthocyanins, a trend of increased concentration of anthocyanins with increased pressure was noted (Figure 86). Anthocyanin degradation was observed at both processing pressures (400 and 550MPa) but was appreciably higher at 400MPa compared with 550 MPa, and these were correlated to antioxidant activity ( $r=0.86$ ) (Del Pozo-Insfran 2007). Anthocyanin aglucones (Figure 87), individual anthocyanins (Table 13), percent anthocyanin polymerization (Figure 88) and total phenolics (Figure 89) were not significantly influenced by high pressure treatments. After 60 min of 600MPa pressure treatment and extreme temperature processing (95°C for 60 min), no changes were observed in the total concentration of lycopene of  $\beta$ -carotene compared to control ( Butz an others 2002). Stability of pigments may be explained by matrix effect: within tissues, the pigments are often compartmentalized and thus protected from adverse influences (Butz an others 2002). Processing pressures of 500 and 800 KPa did not significantly

reduce vitamin C and carotenoid content of treated orange juice (Garcia and others 2001). It has been reported previously that HPP was more detrimental to anthocyanins, ascorbic acid, and color characteristic as compared to thermally pasteurized and control juice's due to oxidase enzymes that were active during HPP processing of muscadine juice (Talcott 2003). Potential mechanisms for destruction include the role of PPO and/or autoxidative mechanisms resulting in co-oxidation of anthocyanins and ascorbic acid (Talcott 2003). In the present study, no PPO activity was detected in the blueberry juice before processing, which can also explain the retention of anthocyanins during high pressure processing. No differences in antioxidant activity measured as disappearance of DPPH over time was noticed (Figure 90). All high pressure treatments significantly increased concentration of the citric acid. Concentration of quinic acid was significantly higher in juice treated at 400MPa than the control and 600MPa treatment (Table 14).




Figure 64. HPLC separation of acid standards detected at 214 nm

4000 ppm quinic, 350 ppm malic, 50 ppm shikimic and 200 ppm citric




Figure 65. HPLC separation of acids detected at 214 nm




Figure 66. Concentration of total anthocyanins after 60 min of maceration as affected by different treatments

Frozen (no maceration), No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 67. Concentration of anthocyanin aglucones after 60 min of maceration as affected by treatments

Frozen (no maceration), No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )

Table 5. Concentration of individual anthocyanins expressed as cyanidin-3-glycoside as affected by different maceration treatments

Anthocyanin	Anthocyanin concentration (ppm)						
	Frozen berries	Control	Benzoate	Heat	LSD	CV	
Delphinidin-3-galactoside	41.60 a	42.25 a	31.42b	29.53b	8.85	12.97	
Delphinidin-3-glucoside	5.49 a	3.94 a	3.04a	5.88a	3.29	38.06	
Cyanidin-3-galactoside	22.49 a	21.29 a	17.45b	17.39a	4.8	12.99	
Delphinidin-3-arabinoside	15.69a	14.14ab	11.64b	10.42b	4.03	16.48	
Cyanidin-3-glucoside	4.49a	3.39a	2.88a	5.00a	2.84	38.32	
Petunidin-3-galactoside	33.01a	33.49a	25.76b	22.52b	5.98	11.08	
Cyanidin-3-arabinoside	8.06a	6.75ab	6.25ab	5.97b	2.18	17.22	
Petunidin-3-glucoside	7.13a	5.04a	4.15a	7.49a	3.77	33.67	
Peonidin-3-galactoside	13.48a	12.68ab	10.44bc	9.03c	2.25	10.49	
Petunidin-3-arabinoside	11.21a	10.32ab	8.63bc	7.24c	2.19	12.47	
Malvidin-3-galactoside	104.26a	100.64a	81.41b	66.60c	13.86	8.34	
Malvidin-3-glucoside	24.05a	16.65ab	14.47b	21.16ab	9.05	25.20	
Malvidin-3-arabinoside	37.21a	35.41a	29.95b	21.5c	4.54	7.78	

abc- Means within the row followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 68. Concentration of total phenolics after 60 min of maceration as affected by different treatment

Frozen (no maceration), No treatment (maceration 55°C 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 69. Changes in polymeric color after 60 min of maceration as affected by different treatment

Frozen (no maceration), No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 70. Activity of PPO ( $\mu\text{mol O}_2/\text{min}/100\text{g}$ ) after 60 min of maceration as affected by different treatment

Frozen (no maceration), No treatment (maceration  $55^\circ\text{C}$ , 1h), Benzoate ( $55^\circ\text{C}$ , 1h, 0.1% sodium benzoate) and Heat ( $90^\circ\text{C}$ , 1min blanching followed by maceration at  $55^\circ\text{C}$  for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )

Table 6 . Concentration of acids in blueberry mash as affected by different maceration treatments

Treatment	Acids mg/100g			
	Quinic	Malic	Shikimic	Citric
Frozen blueberries	941.6 ab	89.4 a	8.9 ab	29.4 b
Control Maceration	868.9 b	79.1 a	9.4 ab	25.7 b
Benzoate Maceration	922.5 ab	103.9 a	7.3 b	25.9 b
Heat Maceration	1072.9 a	71.6 a	15.0 a	42.6 a
LSD	153.4	45.1	6.4	12.7
CV	8.5	27.9	33.2	21.8

ab - Means within the column followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 71. Concentration of total anthocyanins in the pressed juice as affected by different treatment

No treatment (maceration 55°C 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 72. Concentration of anthocyanin aglucones in the pressed juice as affected by different treatment

No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )

Table 7. Concentration of individual anthocyanins expressed as cyanidin-3-glycoside in pressed juice as affected by different maceration treatments

Anthocyanin	Anthocyanin concentration (ppm)				LSD	CV
	Control	Benzoate	Heat			
Delphinidin-3-galactoside	58.71 a	39.62b	63.78a		18.56	17.20
Delphinidin-3-glucoside	7.54a	6.02a	13.43a		8.14	45.28
Cyanidin-3-galactoside	48.51ab	36.64b	57.16a		12.94	13.65
Delphinidin-3-arabinoside	17.96a	11.55a	17.61a		8.86	28.25
Cyanidin-3-glucoside	9.89a	8.04a	16.83a		10.75	46.45
Petunidin-3-galactoside	52.62a	39.72b	56.16a		12.39	12.53
Cyanidin-3-arabinoside	14.07a	10.59a	14.63a		5.30	20.27
Petunidin-3-glucoside	10.17a	9.10a	19.8a		10.70	41.14
Peonidin-3-galactoside	34.34a	28.343a	33.48a		6.79	10.60
Petunidin-3-arabinoside	14.42a	10.48a	13.71a		5.30	20.61
Malvidin-3-galactoside	292.75a	246.93a	263.76a		47.59	8.89
Malvidin-3-glucoside	55.81a	53.97a	86.36a		35.03	26.82
Malvidin-3-arabinoside	91.25a	75.34ba	68.92b		16.78	10.70

140

abc- Means within the raw followed by the same letter are not significantly different (p<0.05)




Figure 73. Concentration of total phenolic in pressed juice expressed as gallic acid equivalents as affected by different maceration treatments

No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 74. Percent polymeric color in pressed juice as affected by different maceration treatments

No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 75. Yield of the pressed juice as affected by different maceration treatments:

No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )

Table 8. Concentration of acids in pressed juice as affected by different maceration treatments

Treatment	Acids mg/1000ml			
	Quinic	Malic	Shikimic	Citric
No treatment	11309.0 a	672.2 a	118.7 a	345.3 b
Benzoate	10914.0 a	728.6 a	113.1 a	256.7 c
Heat	11615.0 a	696.1 a	167.1 a	481.2 a
LSD	3263.0	312.5	86.6	61.8
CV	14.5	23.5	32.6	8.6

abc- Means within the raw followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 76. Concentration of total anthocyanins in pomace as affected by different maceration treatments

No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 77. Concentration of anthocyanin aglucones in pomace as affected by different maceration treatments

No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )

Table 9. Concentration of individual anthocyanins expressed as cyanidin-3-glycoside in pomace as affected by different maceration treatments

Anthocyanin	Anthocyanin concentration (ppm)				
	Control	Benzoate	Heat	LSD	CV
Delphinidin-3-galactoside	88.76 a	83.15a	72.26a	30.19	18.56
Delphinidin-3-glucoside	10.17a	8.89a	13.53a	8.31	38.21
Cyanidin-3-galactoside	44.48a	40.54a	37.9a	15.33	18.73
Delphinidin-3-arabinoside	34.21a	32.03a	28.09a	13.47	21.44
Cyanidin-3-glucoside	8.47a	7.11a	10.46a	7.38	42.56
Petunidin-3-galactoside	70.72a	66.56a	52.53a	21.91	17.34
Cyanidin-3-arabinoside	16.18a	15.19a	14.09a	6.95	22.94
Petunidin-3-glucoside	12.84a	11.36a	16.46a	10.05	37.11
Peonidin-3-galactoside	25.09a	23.36a	18.62a	6.93	15.51
Petunidin-3-arabinoside	24.68a	22.93a	18.34a	8.62	19.63
Malvidin-3-galactoside	194.17a	180.87a	132.59b	46.59	13.78
Malvidin-3-glucoside	37.16a	33.9a	41.30a	20.89	27.82
Malvidin-3-arabinoside	74.81a	68.77a	47.6b	19.21	15.08

abc- Means within the row followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 78. Concentration of total phenolics in pomace expressed as gallic acid equivalents as affected by different maceration treatments

No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 79. Percent polymeric color in pomace as affected by different maceration treatments

No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )

Table 10. Concentration of acids in blueberry pomace as affected by different maceration Treatments

Treatment	Acids mg/100g pomace			
	Quinic	Malic	Shikimic	Citric
No treatment	877.4 a	46.9 b	9.7 ab	nd
Benzoate treated	836.7 a	64.7 a	6.6 b	nd
Heat treated	1032.6 a	55.2 ab	14.3 a	nd
LSD	201.0	14.5	6.9	
CV	10.9	13.1	34.0	

No treatment (maceration 55°C, 1h), Benzoate (55°C, 1h, 0.1% sodium benzoate) and Heat (90°C, 1min blanching followed by maceration at 55°C for 1h). Means followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 80. Concentration of total anthocyanins before and after PEF treatment.

No significant differences were found between treatments ( $p > 0.05$ )

Table 11. Concentration of individual anthocyanins, total phenolic, % polymeric color, total anthocyanins and antioxidant activity before and after PEF treatment

	PEF TREATMENT		LSD	CV
	Before	After		
<b>Individual anthocyanins</b>	Concentration (ppm)			
Delphinidin-3-galactoside	124.42	113.52	24.28	9
Delphinidin-3-glucoside	25.77	23.22	6.12	11.02
Cyanidin-3-galactoside	84.5	78.42	12.52	6.78
Delphinidin-3-arabinoside	44.69	40.19	12.22	12.7
Cyanidin-3-glucoside	26.06	23.61	5.01	8.89
Petunidin-3-galactoside	98.83	91.17	17	7.89
Cyanidin-3-arabinoside	26.95	24.71	6.37	10.88
Petunidin-3-glucoside	32.38	29.59	6.61	9.41
Peonidin-3-galactoside	48.15	44.12	5.83	5.58
Petunidin-3-arabinoside	30.29	27.15	7.56	11.61
Malvidin-3-galactoside	378.83	366.85	67.34	7.97
Malvidin-3-glucoside	119.68	109.61	17.13	6.59
Malvidin-3-arabinoside	124.32	113.56	21.89	8.12
<b>Total Anthocyanins</b>	<b>1164.87</b>	<b>1085.72</b>	<b>198.42</b>	<b>7.77</b>
<b>Other parameters</b>				
% Polymeric color	39.41	38.48	2.28	2.58
Total Phenolics (ppm)	6254.4	5213.7	1134.9	8.73
% DPPH remaining (5min)	31.16	30.71	16.09	22.95

No significant differences ( $p > 0.5$ ) were found between treatments (before and after PEF) in respect to all measured parameters




Figure 81. Spectrophotometric recordings of the disappearance of DPPH at 515 nm in the presence of blueberry juice before and after PEF treatment




Figure 82. Concentration of acids before and after PEF treatment.

No significant differences were found between treatments ( $p > 0.05$ )




Figure 83. Concentration of anthocyanin aglucones before and after pasteurization

No significant differences were found between treatments ( $p > 0.05$ )




Figure 84. Spectrophotometric recordings of the disappearance of DPPH at 515 nm in the presence of blueberry juice before and after pasteurization

Table 12. Concentration of individual anthocyanins, total phenolic, % polymeric color, total anthocyanins and antioxidant activity before and after pasteurization

<b>PASTEURIZATION</b>				
	<b>Before</b>	<b>After</b>	<b>LSD</b>	<b>CV</b>
<b>Individual anthocyanins</b>	Concentration	(ppm)		
Delphinidin-3-galactoside	125.07	123.23	35.26	12.53
Delphinidin-3-glucoside	25.58	25.45	9.18	15.87
Cyanidin-3-galactoside	85.99	85.05	17.11	8.83
Delphinidin-3-arabinoside	45.89	44.67	15.52	15.11
Cyanidin-3-glucoside	26.41	26.42	6.4	10.68
Petunidin-3-galactoside	100.35	98.85	24.19	10.71
Cyanidin-3-arabinoside	27.54	28.31	7.18	11.35
Petunidin-3-glucoside	32.57	32.07	10.66	14.55
Peonidin-3-galactoside	48.76	47.66	7.83	7.16
Petunidin-3-arabinoside	31.16	29.88	9.08	13.13
Malvidin-3-galactoside	397.94	388.69	55.76	6.25
Malvidin-3-glucoside	119.96	117.75	27.51	10.21
Malvidin-3-arabinoside	129.6	124.19	21.65	7.52
<b>Total Anthocyanins</b>	1197.64	1171.44	246.91	9.19
<b>Other parameters</b>				
% Polymeric color	38.78	42.82	4.64	5.02
Total Phenolics (ppm)	4324.8	4713.7	574.3	5.6
% DPPH remaining (5min)	22.22	22.83	7.73	15.15

No significant differences were found between treatments (before and after pasteurization) in respect to all measured parameters




Figure 85. Concentration of acids before and after pasteurization

Means followed by the same letter are not significantly different ( $p \leq 0.05$ )




Figure 86. Concentration of total anthocyanins as affected by different processing pressures

Means followed by the same letter are not significantly different ( $p \leq 0.05$ )




Figure 87. Concentration of anthocyanins aglucones as affected by different processing pressures

No significant differences were found between treatments ( $p > 0.05$ )




Figure 88. Percent anthocyanin polymerization as affected by different processing pressures

Means followed by the same letter are not significantly different ( $p \leq 0.05$ )

Table 13. Concentration of individual anthocyanins expressed as cyanidin-3-glycoside in juice as affected by different processing pressures

Individual Anthocyanins	High Pressure Treatments				LDS	CV
	Control	200 MPa	400 MPa	600 MPa		
Delphinidin-3-galactoside	88.2	83.6	87.1	90.4	25.3	15.4
Delphinidin-3-glucoside	16.9	16.0	16.7	17.3	4.8	15.3
Cyanidin-3-galactoside	66.5	64.0	65.7	67.6	12.7	10.2
Delphinidin-3-arabinoside	29.5	28.0	29.4	30.5	10.5	18.9
Cyanidin-3-glucoside	20.1	19.4	19.8	20.4	3.9	10.5
Petunidin-3-galactoside	73.5	70.0	72.5	75.2	18.3	13.3
Cyanidin-3-arabinoside	20.4	19.3	20.0	20.8	5.4	14.2
Petunidin-3-glucoside	22.7	21.5	22.3	23.2	5.6	13.4
Peonidin-3-galactoside	39.2	37.8	38.8	39.9	6.8	9.3
Petunidin-3-arabinoside	22.0	21.1	22.1	22.7	7.4	17.8
Peonidin-3-glucoside	26.0	25.6	26.1	26.5	3.9	8.1
Malvidin-3-galactoside	299.0	289.0	296.0	303.8	45.3	8.1
Malvidin-3-glucoside	94.3	91.5	93.5	95.7	13.7	7.8
Malvidin-3-arabinoside	102.8	98.6	101.1	104.1	19.0	9.9

156

No significant differences were found between treatments (before and after pasteurization) in respect to all measured parameters




Figure 89. Effect of different processing pressures on total phenolics expresses as ppm of gallic acid equivalents

Means followed by the same letter are not significantly different ( $p \leq 0.05$ )

Table 14. Concentration of acids in juice as affected by High Pressure treatments

High Pressure Treatments						
	Control	200 Mpa	400MPa	600MPa		
<b>Acids</b>	<b>Concentration (ppm)</b>				LDS	CV
Quinic	13,279.0b	13,446.6ab	14,156.5a	13,310.8b	823.5	3.2
Shikimic	181.2ab	179.5b	184.2a	180.4ab	4.0	1.2
Citric	514.7b	614.0a	674.4a	687.2a	84.3	7.2

abc- Means within the raw followed by the same letter are not significantly different ( $p < 0.05$ )




Figure 90. Disappearance of DPPH, at 515 nm, in blueberry juice processed at different pressures for 10 minutes.

## CHAPTER V

### SUMMARY AND CONCLUSIONS

Inhibition of blueberry PPO in food processing is crucial for good quality and the retention of nutrients in products made from blueberries. All inhibitors that were studied decreased the activity of blueberry PPO. Concentration of 0.1 % (maximum allowable for use in food products) and 0.075% of sodium benzoate decreased PPO activity to 5.05% and 8.78%, respectively. Potassium sorbate also decreased PPO activity but almost 50% of activity remained after the maximum allowable concentration of 0.1% was used. Potassium metabisulfite in a concentration as low as 4 ppm inhibited almost 70% of PPO activity. Sulfites (8 ppm) was a strong inhibitor of blueberry PPO which is very important from a practical point of view since amounts of less than 10ppm of sulfites do not have to be declared on the product label (Title 21, U.S. Code of Federal Regulations 101.100).

Fifteen different anthocyanins were identified in the rabbiteye blueberry extract. Derivatives of malvidin were the most abundant (42.1%), followed by delphinidin (19.86%), cyanidin (11.58%), petunidin (16%) and peonidin (6.6 %). Malvidin 3-galactoside was the most abundant individual anthocyanin, with 18.9%. In the model system, anthocyanins were degraded by the action of quinones that were formed by the action of blueberry PPO on 4-methyl catechol. Blueberry PPO degraded 36% of total anthocyanins that were present in the model system. Degradation of anthocyanins was

closely related to their structure. Triphenolic anthocyanins suffered highest percentage of degradation (delphinidin 77 %), followed by diphenolic (petunidin 48% and cyanidin 24%) and monomeric anthocyanins (malvidin 19% and peonidin 16%). Addition of 0.1% sodium benzoate protected 50% of initially degraded anthocyanins by blueberry PPO. Protection of anthocyanin degradation by benzoate was not in the same order for all anthocyanins. The more susceptible that the anthocyanins were to degradation the more they were protected by benzoate. At 0.1 %, benzoate protected approximately 30% of delphinidin, 25% of petunidin, 12% of cyanidin, 13% of malvidin and 8% of peonidin anthocyanins. This is especially important since delphinidin and petunidin anthocyanins have the highest antioxidant activity and impart the characteristic blue color to the blueberries. Although sulfites inhibited PPO activity, they did not protect anthocyanins that were present in the model solution from the action of PPO. Potassium sorbate in the concentration of 0.1% increased the degradation of delphinidin and petunidin anthocyanins.

Not all anthocyanins were equally represented in frozen blueberries. Malvidin glucoside were the most abundant (50.43%) followed by delphinidin (19.3), petunidin (15.64), cyanidin (10.68%) and peonidin (4.1%). Acids identified in frozen blueberry extract were quinic, malic, citric and shikimic. The most abundant acid identified in the blueberry extract was quinic 941.6 mg/100g, followed by malic 89.4 mg/100g, citric 29.4 mg/100g and shikimic 8.9mg/100g.

Maceration treatments before juice extraction influenced PPO activity and phytonutrient content of blueberry mash. Maceration temperature of 55°C reduced PPO activity from 233 to 52.3  $\mu\text{molO}_2/\text{min}/100\text{g}$ . No PPO activity was detected in blueberry

mash that was heat treated at 90°C and then macerated for 60 min at 55°C. Maceration temperature of 55°C was not high enough to affect anthocyanins but on the other hand was high enough to decrease PPO activity almost 4 times thus protecting anthocyanins and red color.

Heat treatment of 90°C that was used to inactivate PPO degraded around 30% of anthocyanins. Although PPO activity can degrade anthocyanins, degradation of anthocyanins in the heat treated (blanched) blueberry mash can be the result of the heat treatment rather than the action of PPO since PPO activity of was barely detectible in the heat treated mash.

Benzoate treatment also significantly reduced total anthocyanins, but possibly due to the action of PPO rather than heat, since 55°C was not enough to degrade anthocyanins. Addition of sodium benzoate decreased PPO activity only around 42% leaving enough active enzyme to degrade anthocyanins.

Not all anthocyanin aglucones were affected the same by the treatments. No significant differences were found between frozen blueberries and control mash in respect to all anthocyanin aglucones. All anthocyanin aglucones except cyanidin aglucones were significantly decreased by heat treatment compared to frozen berries. Heat treatment had the highest negative influence on malvidin anthocyanins (~34 % degraded), followed by delphinidin and petunidin (28% degraded), and cyanidin (20% degraded), in comparison to the frozen berries. Addition of sodium benzoate significantly decreased all anthocyanin aglucones due to the action of PPO that was still active. Treatment with benzoate showed a similar trend of anthocyanin degradation as the heat treatment except that malvidin glycosides were less degraded than with the heat

treatment, with only 25% degradation. These results suggest that malvidin glycosides are more susceptible to heat degradation than to the action of PPO. No significant differences were found between maceration treatments in respect to the total phenolics. Concentrations of quinic acid and citric acids were significantly higher in heat treated mash than in the control maceration treatment. Heat treatment probably helped increase extraction of these acids from the blueberry skin and seeds.

Yield of blueberry juice was around 52% and there were no significant differences between maceration treatments. No significant differences in anthocyanins were found between different maceration treatments in extracted juice. Although heat treated blueberry mash had a significantly lower concentration of total anthocyanins after 60 min of maceration when compared to the control, a lack of differences in anthocyanins in the extracted juice could be the consequence of better extraction of anthocyanins from the heat-broken skin. Recovery of anthocyanins was 12 % from the control and benzoate treated mash, and 17% from the heat treated mash from the frozen fruit.

Malvidin glycosides were the most readily extracted anthocyanins for all treatments, contributing approximately 57-65% of total anthocyanins present in the juice, while their contribution in the frozen blueberry fruit was around 50%. The ratio of delphinidin and petunidin anthocyanins to total anthocyanins decreased in extracted juice compared to their contribution in frozen berries, while cyanidin and peonidin ratios remained the same. Delphinidin anthocyanins decreased the most of all anthocyanins, decreasing from 19% of total anthocyanins present in the frozen berries to 11% in the pressed juice.

Substantial amounts of anthocyanins, 641, 595 and 503 mg/100g remained in the control, benzoate and heat pretreated press cake after the extraction of juice. No significant differences in press cake were found between different treatments in respect to total anthocyanins. Although not significant, heat pretreated press cake had lower amount of anthocyanins than the other two treatments. This can be explained by the fact that the heat pretreatment helped break the skin of the berries which helped the extraction of the anthocyanins into the juice leaving less amounts in the press cake.

Heat extracted juice had a significantly higher amount of total phenolics than benzoate treatment, but not more than the control juice. Less extraction of phenolics from the benzoate treated blueberry mash can be explained by the fact that benzoate treated mash had a lower initial concentration of phenolics. Phenolics percent extraction from the corresponding blueberry mash was 33%, 35% and 41% for the control, benzoate treated and heat treated samples, respectively.

Blueberry mash treatment did not have a significant influence on extracted acids other than citric acid. Heat pretreated extracted juice had significantly higher amounts of citric acid when compared to the control and benzoate pretreated one. Heat treatment probably broke the cells which helped extract more citric acid.

Press cake did not contain any citric acid for any of the treatments indicating that this acid is readily extracted into the juice.

The PEF treatment, kettle pasteurization and HHP treatment did not have a significant influence on total anthocyanins, individual anthocyanins, total phenolics, percent anthocyanin polymerization, and antioxidants in the processed juice. Pasteurization caused a significant increase in quinic and shikimic acids.

From the objectives of this study we can conclude that:

1. Sodium benzoate, potassium metabisulfite and a combination of these two are very good inhibitors of blueberry PPO, while potassium sorbate was the weakest inhibitor.
2. Anthocyanins were protected by sodium benzoate in the model system. Addition of potassium metabisulfite failed to protect anthocyanins and the addition of potassium sorbate increased anthocyanin degradation
3. In respect to inactivation of PPO without negative influence on phytonutrients, especially anthocyanins, maceration treatment for 1h at 55°C would be recommended.
4. Thermal and non thermal processes (PEF and HHP) performed the same in respect to phytonutrients retention.

## REFERENCES

- Adams JB, Ongley MH. 1973. The degradation of anthocyanins in canned strawberries. II. The effect of various additives on the retention of pelargonidin-3-glucoside. *J Food Technol* 8: 305-307.
- Aguilar-Rosas SF, Martin-Belloso O, Ortega-Rivas E, Ballinas-Casarrubias ML, Nevarez-Moorillon GV. 2007. Thermal and pulsed electric fields pasteurization of apple juice: Effects on physicochemical properties and flavour compounds. *Journal Food Eng* 83: 41-46.
- Andriambelason E, Magnier C, Haan-Archipoff G, Lobstein A, Anton R, Beretz A, Stoclet JC, Andriantsitohaina R. 1998. Natural dietary polyphenolic compounds cause endothelium-dependent vasorelaxation in rat thoracic aorta. *J Nutr* 128: 2324-2333.
- Arya SS. 1980. Stability of sorbic acid in aqueous solutions. *J Agric Food Chem* 28: 1246-1249 .
- Bakker J, Bridle P. 1992. Strawberry juice colour: the effect of sulphur dioxide and EDTA on the stability of anthocyanins. *J Sci Food Agric* 60: 477-481.
- Bakker J, Bridle P, Bellworthy SJ, Garcia-Viguera C, Reader HP, Watkins SJ. 1998. Effect of sulphur dioxide and must extraction on colour, phenolic composition and sensory quality of red table wine. *J Sci Food Agric* 78: 297-307.
- Bakker J, Bridle P, Koopman A. 1992. Strawberry juice colour: the effect of some processing variables on the stability of anthocyanins. *J Sci Food Agric* 60: 471-476.
- Ballington JR, Ballinger WE, Maness EP. 1987. Interspecific differences in the percentage of anthocyanins, aglycones, and aglycone-sugars in the fruit of seven species of blueberries. *J Am Soc Hortic Sci* 112: 859-864.
- Barbosa-Canovas GV, Zhang QH, Tabilo-Munizaga G. 2001. Pulsed electric fields in food processing: fundamental aspects and applications / edited by Gustavo V. Barbosa-Canovas, Q. Howard Zhang, Gipsy Tabilo-Munizaga. Food preservation technology series: Lancaster, PA : Technomic Pub. Co., c2001.
- Baruah P, Swain T. 1959. Action of potato phenolase on flavonoid compounds. *J Sci Food Agric* 10: 125-128.

- Billaud C, Lecornu D, Nicolas J. 1996. Substrates and carboxylic acid inhibitors of a partially purified polyphenol oxidase from gum arabic. *J Agric Food Chem* 44: 1668-1675.
- Buchert J, Koponen JM, Suutarinen M, Mustranta A, Lille M, Torronen R, Poutanen K. 2005. Effect of enzyme-aided pressing on anthocyanin yield and profiles in bilberry and blackcurrant juices. *J Sci Food Agric* 85: 2548-2556.
- Bradley D. 2005. Star role for bacteria in controlling flu pandemic? *Nat Rev Drug Discov* 4: 945-946.
- Brand-Williams W, Cuvelier ME, Berset C. 1997. Use of free radical method to evaluate antioxidant activity. *Lebensm. Will. Technol* 30: 609-615
- Campos CA, Alzamora SM, Gerschenson LN. 1997. Sorbate destruction and non-enzymatic browning in model aqueous systems. *Food Sci Technol Int* 3: 405-411.
- Campos CA, Gerschenson LN. 1996. Effect of certain additives on sorbate stability. *Food Res Int* 29: 147-154.
- Carlson JS. 2003. Processing effects on the antioxidant activities of blueberry juices. Raleigh: North Carolina State University. 61 p.
- Casado-Vela J, Selles S, Bru R. 2005. Purification and kinetic characterization of polyphenol oxidase from tomato fruits (*Lycopersicon esculentum* cv. Muchamiel). *J Food Biochem* 29: 381-401.
- Cash JN, Sistrunk WA, Stutte CA. 1976. Characteristics of Concord grape polyphenoloxidase involved in juice color loss. *J Food Sci* 41: 1398-1402.
- Cheyrier V, Souquet JM, Kontek A, Moutounet M. 1994. Anthocyanin degradation in oxidising grape musts. *J Sci Food Agric* 66: 283-288.
- Cortes C, Esteve MJ, Rodrigo D, Torregrosa F, Frigola A. 2006. Changes of colour and carotenoids contents during high intensity pulsed electric field treatment in orange juices. *Food Chem Toxicol* 44: 1932-1939
- Cserhalmi Z, Sass-Kiss A, Toth-Markus M, Lechner N. 2006. Study of pulsed electric field treated citrus juices. *Innovat Food Sci Emerg Tech* 7: 49-54.
- Dallas C, Laureano O. 1994. Effects of pH, sulphur dioxide, alcohol content, temperature and storage time on colour composition of a young Portuguese red table wine. *J Sci Food Agric* 65: 477-485.

- Duckworth HW, Coleman JE. 1970. Physicochemical and kinetic properties of mushroom tyrosinase. *J Biol Chem* 245: 1613-1625.
- Del Pozo-Insfran D, Brenes CH, Talcott ST, Follo-Martinez Ad. 2007. Stability of Copigmented Anthocyanins and Ascorbic Acid in Muscadine Grape Juice Processed by High Hydrostatic Pressure. *J Food Sci* 72: 247-253.
- Deliza R, Rosenthal A, Abadio FBD, Silva CHO, Castillo C. 2005. Application of high pressure technology in the fruit juice processing: benefits perceived by consumers. *J Food Eng* 67: 241-246.
- Dewick PM. 1998. The biosynthesis of shikimate metabolites. *Nat Prod Rep* 15: 17-58.
- Ehlenfeldt MK, Meredith FI, Ballington JR. 1994. Unique organic acid profile of rabbiteye vs. highbush blueberries. *HortScience* : 29: 321-323.
- Elez-Martinez P, Soliva-Fortuny RC, Martinez-Belloso O. 2006. Comparative study on shelf life of orange juice processed by high intensity pulsed electric fields or heat treatment. *Eur Food Res Technol* 222: 321-329.
- Fan X, Mattheis JP. 2001. Inhibition of oxidative and antioxidative enzymes by trans-resveratrol. *J Food Sci* 66: 200-203.
- Ferrar PH, Walker JRL. 1996. Inhibition of diphenol oxidases: a comparative study. *J Biol Chem* 20: 15 30.
- Fernandez Garcia A, Butz P, Bognar A, Tauscher B. 2001. Antioxidative capacity, nutrient content and sensory quality of orange juice and an orange-lemon-carrot juice product after high pressure treatment and storage in different packaging. *Eur Food Res Technol* 213: 290-296.
- Fuleki T, Hope GW. 1964. Effect of various treatments on yield and composition of blueberry juice. *Food Technol April*: 166-168.
- Gandia-Herrero F, Jimenez-Atienzar M, Cabanes J, Garcia-Carmona F, Escribano J. 2005. Differential activation of a latent polyphenol oxidase mediated by sodium dodecyl sulfate. *J Agric Food Chem* 53: 6825-6830.
- Garzon GA, Wrolstad RE. 2002. Comparison of the stability of pelargonidin-based anthocyanins in strawberry juice and concentrate. *J Food Sci* 67: 1288-1299.
- Girelli AM, Mattei E, Messina A, Tarola AM. 2004. Inhibition of polyphenol oxidases activity by various dipeptides. *J Agric Food Chem* 52: 2741 2745.

- Golan-Goldhirsh A, Whitaker JR. 1984. Effect of ascorbic acid, sodium bisulfite, and thiol compounds on mushroom polyphenol oxidase. *J Agric Food Chem* 32: 1003-1009.
- Goodman L, P., Markakis P. 1965. Sulfur dioxide inhibition of anthocyanin degradation by phenolase. *J Food Sci* 30: 135-137.
- Gunata YZ, Sapis JC, Moutounet M. 1987. Substrates and aromatic carboxylic acid inhibitors of grape phenol oxidases. *Phytochemistry* 26: 1573-1575.
- Gil MI, Tomas-Barberan A, Hess-Pierce B, Holcroft DM, Kader AA. 2000. Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. *J Agric Food Chem* 48: 4581-4589.
- Grahl T, Markl H. 1996. Killing of microorganisms by pulsed electric fields. *Appl Microbiol Biotechnol* 45: 148-157.
- Haisman DR. 1974. The effect of sulphur dioxide on oxidising enzyme systems in plant tissues. *J Sci Food Agric* 25: 803-810.
- Ho SY, Mittal GS, Cross JD. 1997. Effects of high field electric pulses on the activity of selected enzymes. *J Food Eng* 31: 69-84.
- Houska M, Strohalm J, Kocurova K, Totusek J, Lefnerova D, Triska J, Vrchotova N, Fiedlerova V, Holasova M, Gabrovska D. 2006. High pressure and foods-fruit/vegetable juices. *J Food Eng* 77: 386-398.
- Iversen CK. 1999. Black currant nectar: Effect of processing and storage on anthocyanin and ascorbic acid content. *J Food Sci* 64: 37-41.
- Indrawati, Van Loey A, Hendrickx M. 2004. Pressure and temperature stability of water-soluble antioxidants in orange and carrot juice: a kinetic study. *Eur food Res Technol* 219: 161-166.
- Janovitz-Klapp AH, Richard FC, Goupy PM, Nicolas JJ. 1990a. Inhibition studies on apple polyphenol oxidase. *J Agric Food Chem* 38: 926-931.
- Janovitz-Klapp AH, Richard FC, Goupy PM, Nicolas JJ. 1990b. Kinetic studies on apple polyphenol oxidase. *J Agric Food Chem* 38: 1437-1441.
- Jin ZT, Zhang QH. 1999. Pulsed electric field inactivation of microorganisms and preservation of quality of cranberry juice. *J Food Process Preserv* 23: 481-497.

- Jensen HD, Krogfelt KA, Cornett C, Hansen SH, Christensen SB. 2002. Hydrophilic carboxylic acids and iridoid glycosides in the juice of American and European cranberries (*Vaccinium macrocarpon* and *V. oxycoccos*), lingonberries (*V. vitis-idaea*), and blueberries (*V. myrtillus*). *J Agric Food Chem* 50: 6871-6874.
- Jurd L. 1963. Reactions involved in sulfite bleaching of anthocyanins. *J Food Sci* 29: 16-19.
- Kader F, Haluk JP, Nicolas JP, Metche M. 1998. Degradation of cyanidin 3-glucoside by blueberry polyphenol oxidase: kinetic studies and mechanisms. *J Agric Food Chem* 46: 3060-3065.
- Kader F, Irmouli M, Nicolas JP, Metche M. 2001. Proposed mechanism for the degradation of pelargonidin 3-glucoside by caffeic acid *o*-quinone. *Food Chem* 75: 139-144.
- Kader F, Irmouli M, Nicolas JP, Metche M. 2002. Involvement of blueberry peroxidase in the mechanisms of anthocyanin degradation in blueberry juice. *J Food Sci* 67: 910-915.
- Kader F, Irmouli M, Zitouni N, Nicolas JP, Metche M. 1999a. Degradation of cyanidin 3-glucoside by caffeic acid *o*-quinone. Determination of the stoichiometry and characterization of the degradation products. *J Agric Food Chem* 47: 4625-4630.
- Kader F, Nicolas JP, Metche M. 1999b. Degradation of pelargonidin 3-glucoside in the presence of chlorogenic acid and blueberry polyphenol oxidase. *J Sci Food Agric* 79: 517-522.
- Kader F, Rovel B, Girardin M, Metche M. 1996. Fractionation and identification of the phenolic compounds of highbush blueberries (*Vaccinium corymbosum*, L.). *Food Chem* 55: 35-40.
- Kader F, Rovel B, Girardin M, Metche M. 1997a. Mechanism of browning in fresh highbush blueberry fruit (*Vaccinium corymbosum* L). Role of blueberry polyphenol oxidase, chlorogenic acid and anthocyanins. *J Sci Food Agric* 74: 31-34.
- Kader F, Rovel B, Girardin M, Metche M. 1997b. Mechanism of browning in fresh highbush blueberry fruit (*Vaccinium corymbosum* L.). Partial purification and characterisation of blueberry polyphenol oxidase. *J Sci Food Agric* 73: 513-516.
- Kahkonen MP, Heinamaki J, Ollilainen V, Heinonen M. 2003. Berry anthocyanins: isolation, identification and antioxidant activities. *J Sci Food Agric* 83: 1403-1411.

- Kalt W, Forney CF, Martin A, Prior RL. 1999. Antioxidant capacity, vitamin C, phenolics, and anthocyanins after fresh storage of small fruits. *J Agric Food Chem* 47: 4638-4644.
- Kalt W, McDonald JE, Donner H. 2000. Anthocyanins, phenolics, and antioxidant capacity of processed lowbush blueberry products. *J Food Sci* 65: 390-393.
- Kim CU, Lew W, Williams MA, Liu H, Zhang L, Swaminathan S, Bischofberger N, Chen MS, Mendel DB, Tai CY, Laver WG, Stevens RC. 1997. Influenza neuraminidase inhibitors possessing a novel hydrophobic interaction in the enzyme active site: design, synthesis, and structural analysis of carbocyclic sialic acid analogues with potent anti-influenza activity. *J Am Chem Soc* 119: 681-690.
- Knorr D. 2003. Impact of non-thermal processing on plant metabolites. *J Food Eng* 56: 131-134.
- Kushman LJ, Ballinger WE. 1968. Acid and sugar changes during ripening in Wolcott blueberries. *Proc Amer Soc Hort Sci* 92: 290-295.
- Lamikanra O, Kirby SD, Musingo MN. 1992. Muscadine grape polyphenoloxidase: partial purification by high pressure liquid chromatography and some properties. *J Food Sci* 57: 686-689.
- Le-Bourvellec C, Le-Quere JM, Sanoner P, Drilleau JF, Guyot S. 2004. Inhibition of apple polyphenol oxidase activity by procyanidins and polyphenol oxidation products. *J Agric Food Chem* 52: 122-130.
- Lee J, Durst RW, Wrolstad RE. 2002. Impact of juice processing on blueberry anthocyanins and polyphenolics: comparison of two pretreatments. *J Food Sci* 67: 1660-1667.
- Lee J, Durst RW, Wrolstad RE. 2005. Determination of total monomeric anthocyanin pigment content of fruit juices, beverages, natural colorants, and wines by the pH differential method: Collaborative study. *J AOAC Int* 88: 1269-1278.
- Lee J, Wrolstad RE. 2004. Extraction of anthocyanins and polyphenolics from blueberry-processing waste. *J Food Sci* 69: 564-573.
- Lila MA. 2004. Anthocyanins and human health: An in vitro investigative approach. *J Biomed Biotechnol* (5):306-313.
- Lu S, Luo Y, Feng H. 2006. Inhibition of apple polyphenol oxidase activity by sodium chlorite. *J Agric Food Chem* 54: 3693-3696.

- Mathew AG, Parpia HAB. 1971. Food browning as a polyphenol reaction. *Adv Food Res* 19: 75-145.
- Mathew AG, Parpia HAB. 1971. Food browning as a polyphenol reaction. *Adv Food Res* 19: 75-145.
- Mayer AM. 1987. Polyphenol oxidases in plants--recent progress. *Phytochemistry* 26: 11-20.
- Mayer AM. 2006. Polyphenol oxidases in plants and fungi: Going places? A review. *Phytochemistry* 67: 2318-2331.
- Mayer AM, Harel E. 1979. Polyphenol oxidases in plants. *Phytochemistry* 18: 193-215.
- Mazza G, Brouillard R. 1987a. Color stability and structural transformations of cyanidin 3,5-diglucoside and four 3-deoxyanthocyanins in aqueous solutions. *J Agric Food Chem* 35: 422-426.
- Mazza G, Brouillard R. 1987b. Recent developments in the stabilization of anthocyanins in food products. *Food Chem* 25: 207-225.
- Mazza G, Miniati E. 1993. Anthocyanins in fruits, vegetables, and grains / G. Mazza, E. Miniati. *Anthocyanins in fruits, vegetables, and grains* / G. Mazza, E. Miniati. Boca Raton : CRC Press.
- Main G, Faupel M, Morris J, McNew R. 2001. Quality and stability of blueberry juice blended with apple, grape and cranberry juice. *J Food Qual* 24: 111-125.
- Markakis PA, Jarczyk A, Krishna SP. 1963. Nonvolatile acids of blueberries. *J Agric Food Chem* 11: 8-11.
- McGhie TK, Ainge GD, Barnett LE, Cooney JM, Jensen DJ. 2003. Anthocyanin glycosides from berry fruit are absorbed and excreted unmetabolized by both humans and rats. *J Agric Food Chem* 51: 4539-4548.
- Montgomery MW, Reyes FGR, Cornwell C, Beavers DV. 1982. Sugars and acid analysis and effect of heating on color stability of Northwest Concord grape juice. *J Food Sci* 47: 1883.
- Moyer RA, Hummer KE, Finn CE, Frei B, Wrolstad R. 2002. Anthocyanins, phenolics, and antioxidant capacity in diverse small fruits: *Vaccinium*, *Rubus*, and *Ribes*. *J Agric Food Chem* 50: 519-525.

- Murata M, Tsurutani M, Tomita M, Homma S, Kaneko K. 1995. Relationship between apple ripening and browning: changes in polyphenol content and polyphenol oxidase. *J Agric Food Chem* 43: 1115-1121.
- Ofek I, Goldhar J, Sharon N. 1996. Anti-*Escherichia coli* adhesin activity of cranberry and blueberry juices. *Anti-Escherichia coli* adhesin activity of cranberry and blueberry juices. Toward anti-adhesion therapy for microbial diseases: New York : Plenum Press, c1996. p 179-83.
- Nakajima J, Tanaka I, Seo S, Yamazaki M, Saito K. 2004. LC/PDA/ESI-MS profiling and radical scavenging activity of anthocyanins in various berries. *J Biomed Biotechnol* (5): 241-247.
- Nicolas J, Potus J. 1994. Enzymatic oxidation phenomena and coupled oxidations. Effects of lipoxygenase in breadmaking and of polyphenol oxidase in fruit technology. *Sci Aliments* 14: 627-642.
- Nunez-Delicado E, Serrano-Megias M, Perez-Lopez AJ, Lopez-Nicolas JM. 2005. Polyphenol oxidase from Dominga table grape. *J Agric Food Chem* 53: 6087-6093.
- Pansare SV, Adsool VA. 2006. A concise, enantioselective approach to (-)-quinic acid. *Org Lett* 8 : 2035-2037.
- Peng C, Y., Markakis P. 1963. Effect of phenolase on anthocyanins. *Nature* 199: 597-598.
- Pifferi PG, Baldassarii L, Cultrera R. 1974. Inhibition by carboxylic acids of an *o*-diphenol oxidase from *Prunus avium* fruits. *J Sci Food Agric* 25: 263-270.
- Pifferi PG, Cultrera R. 1974. Enzymatic degradation of anthocyanins: the role of sweet cherry polyphenol oxidase. *J Food Sci* 39: 786-791.
- Prior RL, Lazarus SA, Cao G, Muccitelli H, Hammerstone JF. 2001. Identification of procyanidins and anthocyanins in blueberries and cranberries (*Vaccinium* spp.) using high-performance liquid chromatography/mass spectrometry. *J Agric Food Chem* 49: 1270-1276.
- Prior RL, Ehlenfeldt M, Lischner N, Kalt W, Mainland CM, Krewer G, Martin A, Cao G, Sofic E, O'Brien C, McEwen J. 1998. Antioxidant capacity as influenced by total phenolic and anthocyanin content, maturity, and variety of *Vaccinium* species. *J Agric Food Chem* 46: 2686-2693.
- Raynal J, Moutounet M. 1989. Intervention of phenolic compounds in plum technology. 2. Mechanisms of anthocyanin degradation. *J Agric Food Chem* 37: 1051-1053.

- Ribereau-Gayon P, Pontallier P, Glories YO. 1983. Some interpretations of colour changes in young red wines during their conservation Anthocyanins, tannins. *J Sci Food Agric* 34: 506-516.
- Rein MJ, Heinonen M. 2004. Stability and enhancement of berry juice color. *J Agric Food Chem* 52: 3106-3114.
- Rossi M, Giussani E, Morelli R, Scalzo RL, Nani RC, Torreggiani D. 2003. Effect of fruit blanching on phenolics and radical scavenging activity of highbush blueberry juice. *Food Res Intern* 36: 999-1005.
- Roberts JS, Gentry TS, Bates AW. 2004. Utilization of dried apple pomace as a press aid to improve the quality of strawberry, raspberry, and blueberry juices. *J Food Sci* 69: 181-190.
- Sakamura S, Obata Y. 1963. Anthocyanase and anthocyanins in eggplant, *Solanum melongena* L. Part II. Isolation and identification of chlorogenic acid and related compounds from eggplant. *Agric Biol Chem* 27: 121-127.
- Sakamura S, Watanase S, Obata Y. 1965. Oxidative decolorization of the anthocyanin by polyphenol oxidase. *Agric Biol Chem* 29: 181-190.
- Sanchez-Ferrer A, Bru R, Valero E, Garcia-Carmona F. 1989. Changes in pH-dependent grape polyphenoloxidase activity during maturation. *J Agric Food Chem* 37: 1242-1245.
- Sapers GM. 1993. Browning of foods: control by sulfites, antioxidants, and other means. *Food Technol* 47: 75-84.
- Sapers GM, Burgher AM, Phillips JG, Jones SB, Stone EG. 1984. Color and composition of highbush blueberry cultivars. *J Am Soc Hortic Sci* 109: 105-111.
- Sapers GM, Hicks KB, Phillips JG, Garzarella L, Pondish DL, Matulaitis RM, McCormack TJ, Sondey SM, Seib PA, El-Atawy YS. 1989. Control of enzymatic browning in apple with ascorbic acid derivatives, polyphenol oxidase inhibitors, and complexing agents. *J Food Sci* 54: 997-1002.
- Sarni-Manchado P, Cheynier V, Moutounet M. 1997. Reactions of polyphenoloxidase generated caftaric acid *o*-quinone with malvidin 3-*O*-glucoside. *Phytochemistry* 45: 1365-1369.
- Sarni P, Fulcrand H, Souillol V, Souquet J, Cheynier V. 1995. Mechanism of Anthocyanin Degradation in Grape Must-Like Model Solution. *J Sci Food Agric* 69: 385-391.

- Sayavedra-Soto LA, Montgomery MW. 1986. Inhibition of polyphenoloxidase by sulfite. *J Food Sci* 51: 1531-1536.
- Schmidt BM, Erdman JW, Jr., Lila MA. 2005. Effects of food processing on blueberry antiproliferation and antioxidant activity. *J Food Sci* 70: 389-394.
- Senorans FJ, Ibanez E, Cifuentes A. 2003. New trends in food processing. *Crit Rev Food Sci Nutr* 43: 507-526.
- Sellappan S, Akoh CC, Krewer G. 2002. Phenolic compounds and antioxidant capacity of Georgia-grown blueberries and blackberries. *J Agric Food Chem* 50: 2432-2438.
- Serradell Md-I-A, Rozenfeld PA, Martinez GA, Civello PM, Chaves AR, Anon MC. 2000. Polyphenoloxidase activity from strawberry fruit (*Fragaria x ananassa*, Duch., cv Selva): characterisation and partial purification. *J Sci Food Agric* 80: 1421-1427.
- Shahidi F, Naczki M. 1995. Food phenolics: Sources, Chemistry, Effects and Applications. Technomic Publishing Company, Lancaster PA.
- Siddiq M, Sinha NK, Cash JN. 1992. Characterization of polyphenoloxidase from Stanley plums. *J Food Sci* 57: 1177-1179.
- Sistrunk WA, Morris JR, Kozup J. 1982. The effect of chemical treatments and heat on color stability of frozen machine-harvested strawberries for jam. *J Am Soc Hortic Sci* 107: 693-697.
- Skrede G, Wrolstad RE, Durst RW. 2000. Changes in anthocyanins and polyphenolics during juice processing of highbush blueberries (*Vaccinium corymbosum* L.). *J Food Sci* 65: 357-364.
- Sojo MM, Nunez-Delgado E, Garcia-Carmona F, Sanchez-Ferrer A. 1999. Cyclodextrins as activator and inhibitor of latent banana pulp polyphenol oxidase. *J Agric Food Chem* 47: 518-523.
- Soliva-Fortuny RC, Elez-Martinez P, Sebastian-Caldero M, Martin-Belloso O. 2002. Kinetics of polyphenol oxidase activity inhibition and browning of avocado puree preserved by combined methods. *J Food Eng* 55: 131-137.
- Talcott ST, Brenes CH, Pires DM, Del Pozo-Insfran D. 2003. Phytochemical stability and color retention of copigmented and processed muscadine grape juice. *J Agric Food Chem* 51: 957-963.

- Timberlake C, F., Bridle P. 1967a. Flavylum salts, anthocyanidins and anthocyanins. Reactions with sulphur dioxide. *J Sci Food Agric* 18: 479-485.
- Timberlake C, F., Bridle P. 1967b. Flavylum salts, anthocyanidins and anthocyanins. Structural transformations in acid solutions. *J Sci Food Agric* 18: 473-478.
- Timberlake CF, Bridle P. 1971. The anthocyanins of apples and pears: the occurrence of acyl derivatives, 22 (10); 509-513.
- Threlfall RT, Morris JR, Howard LR, Brownmiller CR, Walker TR. 2005. Pressing effects on yield, quality, and nutraceutical content of juice, seeds, and skins from black beauty and sunbelt grapes. *J Food Sci* 70: 167-171.
- Tomas-Barberan FA, Espin JC. 2001a. Phenolic compounds and related enzymes as determinants of quality in fruits and vegetables. *J Sci Food Agric* 81: 853-876.
- Tomas-Barberan FA, Espin JC. 2001b. Phenolic compounds and related enzymes as determinants of quality in fruits and vegetables. *J Sci Food Agric* 81: 853-876.
- Toepfl S, Mathys A, Heinz V, Knorr D. 2006. Review: Potential of high hydrostatic pressure and pulsed electric fields for energy efficient and environmentally friendly food processing. *Food Rev Int* 22: 405-423.
- Tsuda T, Horio F, Uchida K, Aoki H, Osawa T. 2003. Dietary cyanidin 3-O-beta-D-glucoside-rich purple corn color prevents obesity and ameliorates hyperglycemia in mice. *J Nutr* 133: 2125-2130.
- Unal MU, Sener A. 2006. Determination of some biochemical properties of polyphenol oxidase from Emir grape (*Vitis vinifera* L. cv. Emir). *J Sci Food Agric* 86: 2374-2379.
- Valero E, Varon R, Garcia-Carmona F. 1992. Kinetic study of the effect of metabisulfite on polyphenol oxidase. *J Agric Food Chem* 40: 904-908.
- Vamos-Vigyazo L. 1981. Polyphenol oxidase and peroxidase in fruits and vegetables. *CRC Crit Rev Food Sci Nutr* 15: 49-127.
- Walker JRL, McCallion RF. 1980. The selective inhibition of *ortho*- and *para*-diphenol oxidases. *Phytochemistry* 19: 373-377.
- Walker JRL, Wilson EL. 1975. Studies on the enzymic browning of apples. Inhibition of apple *o*-diphenol oxidase by phenolic acids. *J Sci Food Agric* 26: 1825-1831.
- Wang H, Prior RL, Cao G. 1997. Oxygen radical absorbing capacity of anthocyanins. *J Agric Food Chem* 45: 304-309.

- Wedzicha BL. 1992. Chemistry of sulphiting agents in food. *Food Addit Contam* 9: 449-459.
- Wedzicha BL, Bellion I, Goddard SJ. 1991. Inhibition of browning by sulfites. *Adv Exp Med Biol* 289: 217-236.
- Wesche-Ebeling P, Montgomery MW. 1990a. Strawberry polyphenoloxidase: extraction and partial characterization. *J Food Sci* 55(5): 1320-1324.
- Wesche-Ebeling P, Montgomery MW. 1990b. Strawberry polyphenoloxidase: its role in anthocyanin degradation. *J Food Sci* 55(3): 731-734.
- Wesche-Ebeling P, Montgomery MW. 1990c. Strawberry polyphenoloxidase: purification and characterization. *J Food Sci* 55(5): 1315-1319.
- Weemaes CA, Cordt Svd, Ludikhuyze LR, Broeck Ivd, Hendrickx ME, Tobback PP. 1997. Influence of pH, benzoic acid, EDTA, and glutathione on the pressure and/or temperature inactivation kinetics of mushroom polyphenoloxidase. *Biotechnol Progr* 13: 25-32.
- Weemaes CA, Ludikhuyze LR, Broeck Ivd, Hendrickx ME, Tobback PP. 1998. Activity, electrophoretic characteristics and heat inactivation of polyphenoloxidases from apples, avocados, grapes, pears and plums. *Lebensm Wiss Technol Food Sci Technol* 31: 44-49.
- Wesierska E, Trziszka T. 2007. Evaluation of the use of pulsed electrical field as a factor with antimicrobial activity. *J Food Eng*; 78: 1320-1325.
- Wouters PC, Alvarez I, Raso J. 2001. Critical factors determining inactivation kinetics by pulsed electric field food processing. *Trends Food Sci Technol* 12: 112-121.
- Wrolstad RE. 2001. *Current protocols in food analytical chemistry*. RE Wrolstad (Ed.) New York: John Willey & Sons.
- Wrolstad RE. 1976. Color and pigment analyses in fruit products. Bul 624. Oregon State University Agriculture Experiment Station, Corvallis OR.
- Wrolstad RE. 2004. Anthocyanin pigments-bioactivity and coloring properties. *J Food Sci* 69: 419-421.
- Wrolstad RE, Lee DD, Poesi MS. 1980. Effect of microwave blanching on the color and composition of strawberry concentrate. *J Food Sci* 45: 1573-1577.

- Wrolstad RE, Wightman JD, Durst RW. 1994. Glycosidase activity of enzyme preparations used in fruit juice processing. *Food Technol* 48:90.
- Yang CST, Yang PPA. 1987. Effect of pH, certain chemicals and holding time-temperature on the color of lowbush blueberry puree. *J Food Sci* 52: 346.
- Yang RJ, Li SQ, Zhang QH. 2004. Effects of pulsed electric fields on the activity of enzymes in aqueous solution. *J Food Sci* 69: 241-248
- Yemenicioglu A, Ozkan M, Cemeroglu B. 1997. Heat inactivation kinetics of apple polyphenoloxidase and activation of its latent form. *J Food Sci* 62: 508-510.
- Yeom HW, McCann KT, Streaker CB, Zhang QH. 2002. Pulsed electric field processing of high acid liquid foods: a review. *Adv Food Nutr Res* 44: 32-44.
- Yokotsuka K, Singleton VL. 1997. Disappearance of anthocyanins as grape juice is prepared and oxidized with PPO and PPO substrates. *Am J Enol Viticult* 48: 13-25.
- Zafrilla P, Ferreres F, Tomas-Barberan FA. 2001. Effect of processing and storage on the antioxidant ellagic acid derivatives and flavonoids of red raspberry (*Rubus idaeus*) jams. *J Agric Food Chem* 49: 3651-3655.
- Zhong K, Chen F, Wu JH, Wang ZF, Liao XJ, Hu XS, Zhang ZH. 2005. Kinetics of inactivation of *Escherichia coli* in carrot juice by pulsed electric field. *J Food Process Eng* 28: 595-609.
- Zhang Y, Liao X, Ni Y, Wu J, Hu X, Wang Z, Chen F. 2007. Kinetic analysis of the degradation and its color change of cyanidin-3-glucoside exposed to pulsed electric field. *Z Lebensm Unters F A* 224: 597-603.
- Zuckerman H, Krasik YE, Felsteiner J. 2004. Effect of pulsed shock waves on the activity of polyphenol oxidase, alkaline phosphatase and green fluorescent protein. *J Sci Food Agric* 84: 841-844.